

f -cohomology and motives over number rings

Jakob Scholbach *

March 5, 2010

Abstract

This paper is concerned with an interpretation of f -cohomology, a modification of motivic cohomology of motives over number fields, in terms of motives over number rings. Under standard assumptions on mixed motives over finite fields, number fields and number rings, we show that the two extant definitions of f -cohomology of mixed motives M_η over F —one via ramification conditions on ℓ -adic realizations, another one via the K -theory of proper regular models—both agree with motivic cohomology of $\eta_{l*}M_\eta[1]$. Here η_{l*} is constructed by a limiting process in terms of intermediate extension functors j_{l*} defined in analogy to perverse sheaves.

The aim of this paper is to give an interpretation of f -cohomology in terms of motives over number rings. The notion of f -cohomology goes back to Beilinson who used it to formulate a conjecture about special L -values [Beĭ84, Beĭ86]. The most classical example is what is now called $H_f^1(F, \mathbf{1}(1))$, f -cohomology of $\mathbf{1}(1)$, the motive of a number field F , twisted by one. This group is $\mathcal{O}_F^\times \otimes_{\mathbb{Z}} \mathbb{Q}$, as opposed to the full motivic cohomology $H^1(F, \mathbf{1}(1)) = F^\times \otimes \mathbb{Q}$. Together with the Dirichlet regulator, it explains the residue of the Dedekind zeta function $\zeta_F(s)$ at $s = 1$. In general, for any mixed motive M_η over F , there are two independent yet conjecturally equivalent ways to define $H_f^1(F, M_\eta) \subset H^1(F, M_\eta)$. We propose a third way to obtain these groups, by interpreting f -cohomology as motivic cohomology of suitable motives over $\mathcal{O}_F$. This idea is due to Huber.

Let us briefly recall the two definitions of $H_f^1(M_\eta)$. The first is due to Beilinson [Beĭ87a, Remark 4.0.1.b], Bloch and Kato [BK90, Conj. 5.3.] and Fontaine [Fon92, FPR94]. It is given by picking elements in motivic cohomology acted on by the local Galois groups in a prescribed way (Definition 6.1, Definition 6.4, Definition 6.6). The second definition of $H_f^1(M_\eta)$, due to Beilinson [Beĭ86, Section 8], applies to $M_\eta = h^{i-1}(X_\eta)(n)$, with X_η smooth and projective over F , $i - 2n < 0$. It is given by the image of K -theory of a regular proper model X of X_η (Definition 6.11). Such a model may not exist, but there is a unique meaningful extension of this definition to all Chow motives over F due to Scholl [Sch00].

*Universität Münster, Mathematisches Institut, Einsteinstr. 62, D-48149 Münster, Germany, jakob.scholbach@uni-muenster.de

Our main results (Theorems 6.9, 6.12, 6.14) show that both definitions of $H_F^1(M_\eta)$ agree with $H^0(\eta_{!*}h^{i-1}(X_\eta, n)[1])$. Here $\eta_{!*}$ is a functor that attaches to any suitable mixed motive over F one over $\mathcal{O}_F$. It is defined by a limiting process using the intermediate extension $j_{!*}$ familiar from perverse sheaves [BBD82] along open immersions $j : U \rightarrow \text{Spec } \mathcal{O}_F$. Even to formulate such a definition, one has to rely on profound conjectures, namely the existence of mixed motives over (open subschemes of) $\text{Spec } \mathcal{O}_F$. The proof of the two main theorems also requires us to assume a number of properties related to weights of motives and, for the first comparison result, Beilinson’s conjecture about the agreement of rational and numerical equivalence on smooth projective varieties over finite fields up to torsion (Conjecture 6.7).

A motivation for the results of this work lies in an application to special L -values conjectures [Sch10]. Very briefly, Beilinson’s conjecture concerning special L -values for mixed motives M_η over $\mathbb{Q}$ has f -cohomology as motivic input. L -functions of such motives can be generalized to motives over $\mathbb{Z}$ such that the classical L -function of M_η agrees with the L -function (over $\mathbb{Z}$) of $\eta_{!*}M_\eta[1]$. Thereby the L -function and the motivic data in Beilinson’s conjecture belong to the same motive over $\mathbb{Z}$, thus giving content to a general conjecture about special L -values for motives over $\mathbb{Z}$. In this light it is noteworthy that $H^0(\eta_{!*}h^{2n-1}(X_\eta, n)[1])$ identifies with the group that occurs in the part of Beilinson’s conjecture that describes special values at the central point.

The contents of the paper are as follows: Section 1 is the basis of the remainder; it lists a number of axioms on triangulated categories of motives. Such categories $\mathbf{DM}_{\text{gm}}(S)$ have been constructed by Voevodsky [Voe00] and Hanamura [Han95] (over fields) and Levine [Lev98] (over bases S over a field). The various approaches are known to be (anti-)equivalent, at least for rational coefficients [Lev98, Section VI.2.5], [Bon09, Section 4]. Over more general bases S , a candidate for the category $\mathbf{DM}(S)$ has been constructed by Ivorra [Ivo07]. Cisinski and Déglise are developing a robust theory of such motives [CD10]. We sum up the properties of this construction by specifying a number of axioms concerning triangulated categories of motives that will be used in the sequel. They are concerned with the “core” behavior of $\mathbf{DM}(S)$, that is: functoriality, compactness, the monoidal structure and the relation to algebraic K -theory, as well as localization, purity, base-change and resolution of singularities. We work with motives with rational coefficients only, since this is sufficient for all our purposes. We use a contravariant notation for motives, that is to say the functor that maps any scheme X to its motive $M(X)$ shall be contravariant. This is in line with most pre-Voevodsky papers.

Section 2 is a very brief reminder on realizations. The existence of various realizations, due to Huber and Ivorra [Hub95, Hub00, Ivo07], is pinning down the intuition that motives should be universal among (reasonable) cohomology theories.

Section 4 spells out a number of conjectural properties (also called axioms in the sequel) of $\mathbf{DM}_{\text{gm}}(S)$, where S is either a finite field $\mathbb{F}_p$, a number field F or a number ring $\mathcal{O}_F$. The first group of these properties centers around the existence of a category of mixed motives $\mathbf{MM}(S)$, which is to be the heart of the so-called

motivic t -structure. The link between mixed motives over $\mathcal{O}_F$ and $\mathbb{F}_p$ or F is axiomatized by mimicking the exactness properties familiar from perverse sheaves (Axiom 4.2). On the triangulated subcategory $\mathbf{DATM}(\mathcal{O}_F) \subset \mathbf{DM}_{\text{gm}}(\mathcal{O}_F)$ of Artin-Tate motives, a t -structure satisfying these properties is constructed in a separate paper [Scha]. A key requirement on mixed motives is that the realization functors on motives should be exact (Axiom 4.8). For the ℓ -adic realization over $\text{Spec } \mathcal{O}_F[1/\ell]$, this requires a notion of perverse sheaves over that base, which is provided in Section 3. Another important conjectural facet of mixed motives are weights. Weights are an additional structure encountered in both Hodge structures and ℓ -adic cohomology of algebraic varieties over finite fields, both due to Deligne [Del74, Del80]. They are important in that morphisms between Hodge structures or ℓ -adic cohomology groups are known to be strictly compatible with weights, moreover, they are respected to a certain extent by smooth maps and proper maps. It is commonly assumed that this should be the case for mixed motives, too.

The remaining two sections assume the validity of the axiomatic framework set up so far. The first key notion in Section 5 is the intermediate extension $j_{!*}M$ of a mixed motive M along some open embedding j inside $\text{Spec } \mathcal{O}_F$. This is done in strict parallelity to the case of perverse sheaves, due to Beilinson, Bernstein and Deligne [BBD82]. Quite generally, much of this paper is built on the idea that the abstract properties of mixed perverse sheaves (should) give a good model for mixed motives over number rings. Next we develop a notion of smooth motives, which is an analog of lisse étale sheaves. This is needed to extend the intermediate extension to an extension functor $\eta_{!*}$ that extends motives over F to ones over $\mathcal{O}_F$. Finally, we apply the axiom on the exactness of ℓ -adic realization to show that intermediate extensions commute with the realization functors. This will be a stepstone in a separate work on L -functions of motives [Sch10].

Section 6 gives the comparison theorems on f -cohomology mentioned above. The two definitions of f -cohomology being quite different, the proofs of the comparison statements are different, too: the first is essentially based on the Hochschild-Serre spectral sequence. The crystalline case of that definition of f -cohomology is disregarded throughout. The second proof is a purely formal, if occasionally intricate bookkeeping of cohomological degrees and weights.

The problem of finding a motivic interpretation of terms such as $H_f^1(M_\eta)$ underlying the formulation of Beilinson's conjecture has been studied by Scholl [Sch91, Sch00, Schb], who develops an abelian category $\mathbf{MM}(F/\mathcal{O}_F)$ of mixed motives over $\mathcal{O}_F$ by taking mixed motives over F , and imposing additional non-ramification conditions. Conjecturally, the group $\text{Ext}_{\mathbf{MM}(F/\mathcal{O}_F)}^i(\mathbf{1}, h^i(X_\eta, n))$ for X_η/F smooth and projective, $i = 0, 1$, agrees with what amounts to $H^{i-1}(\eta_{!*}h^{2n-1}(X_\eta, n)[1])$.

No originality is claimed for Sections 1, 2, and 4, except perhaps for the formulation of the relation of mixed motives over $\mathcal{O}_F$ and F and the residue fields $\mathbb{F}_p$, which however is a natural and immediate translation of the theory of perverse sheaves. I would like to thank Denis-Charles Cisinski and Frédéric Déglise for communicating to me their work on $\mathbf{DM}_{\text{gm}}(S)$ over general bases

[CD10]. Finally, I gratefully acknowledge Annette Huber's advice in writing my thesis, of which this paper is a part.

1 Geometric motives

Throughout this paper, F is a number field, $\mathcal{O}_F$ its ring of integers, $\mathfrak{p}$ stands for a place of F . For finite places, the residue field is denoted $\mathbb{F}_{\mathfrak{p}}$. By scheme we mean a Noetherian separated scheme. Actually, it suffices to think of schemes of finite type over one of the rings just mentioned. In this section S denotes a fixed base scheme.

This section is setting up a number of axioms describing a triangulated category $\mathbf{DM}_{\text{gm}}(S)$ of geometric motives over S . They will be used throughout this work. As pointed out in the introduction, the material of this section is due to Cisinski and Déglise [CD10], who build such a category of motives using Ayoub's base change formalism [Ayo07].

Axiom 1.1. (Motivic complexes and functoriality)

- There is a triangulated $\mathbb{Q}$ -linear category $\mathbf{DM}(S)$. It is called category of *motivic complexes* over S (with rational coefficients). It has all limits and colimits.
- (Tensor structure) The category $\mathbf{DM}(S)$ is a triangulated symmetric monoidal category (see e.g. [Lev98, Part 2, II.2.1.3]). Tensor products commute with direct sums. The unit of the tensor structure is denoted $\mathbf{1}_S$ or $\mathbf{1}$. Also, there are internal Hom-objects in $\mathbf{DM}$, denoted $\underline{\text{Hom}}$. The dual $M^\vee$ of an object $M \in \mathbf{DM}(S)$ is defined by $M^\vee := \underline{\text{Hom}}(M, \mathbf{1})$.
- For any map $f : X \rightarrow Y$ of schemes, there are pairs of adjoint functors

$$f^* : \mathbf{DM}(Y) \rightleftarrows \mathbf{DM}(X) : f_*$$

such that $f^*\mathbf{1}_Y = \mathbf{1}_X$ and, if f is quasi-projective,

$$f_! : \mathbf{DM}(X) \rightleftarrows \mathbf{DM}(Y) : f^!$$

The existence of $f_!$ and $f^!$ is restricted to quasi-projective maps since the abstract construction of these functors in Ayoub's work [Ayo07, Section 1.6.5], on which Cisinski's and Déglise's construction of motives over general bases [CD10] relies, has a similar restriction.

Recall that an object X in a triangulated category $\mathcal{T}$ closed under arbitrary direct sums is *compact* if $\text{Hom}(X, -)$ commutes with direct sums. The subcategory of $\mathcal{T}$ of compact objects is triangulated and closed under direct summands (a.k.a. a *thick subcategory*) [Nee01, Lemma 4.2.4]. The category $\mathcal{T}$ is called *compactly generated* if the smallest triangulated subcategory closed under arbitrary sums containing the compact objects is the whole category $\mathcal{T}$.

Axiom 1.2. (Compact objects) The motive $\mathbf{1} \in \mathbf{DM}(S)$ is compact. The functors f^* and $f^!$, whenever defined, and $\otimes$ and $\underline{\mathbf{Hom}}$ preserve compact objects. The same is true for f_* and $f_!$ if f is of finite type. The canonical map $M \rightarrow (M^\vee)^\vee$ is an isomorphism for any compact object M .

Definition 1.3. The subcategory of compact objects of $\mathbf{DM}(S)$ is denoted $\mathbf{DM}_{\text{gm}}(S)$ and called the category of *geometric motives* over S .

For any map $f : X \rightarrow S$ of finite type, the object $M_S(X) := M(X) := f_* f^* \mathbf{1} \in \mathbf{DM}_{\text{gm}}(S)$ is called the *motive* of X over S . By adjunction, M is a contravariant functor from schemes of finite type over S to $\mathbf{DM}_{\text{gm}}(S)$. For any quasi-projective $f : X \rightarrow S$, the *motive with compact support* of X , $M_c(X)$, is defined as $f_! f^* \mathbf{1} \in \mathbf{DM}_{\text{gm}}(S)$.

The smallest thick subcategory of $\mathbf{DM}(S)$ containing the image of M is denoted $\mathbf{DM}_{\text{gm}}^{\text{eff}}(S)$ and called the category of *effective geometric motives*. The closure of that subcategory under all direct sums is called the category of *effective motives*, $\mathbf{DM}_{\text{eff}}(S)$.

Axiom 1.4. (Tensor product vs. fiber product) The functor M is an additive tensor functor, i.e., maps disjoint unions of schemes over S to direct sums and fiber products of schemes over S to tensor products in $\mathbf{DM}_{\text{gm}}(S)$.

Axiom 1.5. (Compact generation) The categories $\mathbf{DM}(S)$ and $\mathbf{DM}_{\text{eff}}(S)$ are compactly generated.

There is a decomposition $M(\mathbb{P}_S^1) = \mathbf{1} \oplus \mathbf{1}(-1)[-2]$, where

$$\mathbf{1}(-1) := \ker(M(\mathbb{P}_S^1) \rightarrow M(S) \rightarrow M(\mathbb{P}_S^1)[2]),$$

where the first map is induced by the projection onto the base, the second map stems from the rational point $0 \in \mathbb{P}_S^1$. The object $\mathbf{1}(-1)$ is called *Tate object* or *Tate motive*. The category $\mathbf{DM}(S)$, being closed under countable direct sums is pseudo-abelian [Lev98, Lemma II.2.2.4.8.1], i.e., it contains kernels of projectors. It follows from the preceding axioms that $\mathbf{1}(-1) \in \mathbf{DM}_{\text{gm}}^{\text{eff}}(S)$.

Axiom 1.6. (Cancellation and Effectivity) In $\mathbf{DM}_{\text{gm}}(S)$ (and thus in $\mathbf{DM}(S)$), the Tate object $\mathbf{1}(-1)$ has a tensor-inverse denoted $\mathbf{1}(1)$. For any $M \in \mathbf{DM}(S)$, $n \in \mathbb{Z}$, set $M(n) := M \otimes \mathbf{1}(1)^{\otimes n}$. Then there is a canonical isomorphism called *cancellation* isomorphism ($n \in \mathbb{Z}$, $M, N \in \mathbf{DM}(S)$):

$$\text{Hom}_{\mathbf{DM}(S)}(M, N) \cong \text{Hom}_{\mathbf{DM}(S)}(M(n), N(n)).$$

The smallest tensor subcategory of $\mathbf{DM}_{\text{gm}}(S)$ that contains $\mathbf{DM}_{\text{gm}}^{\text{eff}}(S)$ and $\mathbf{1}(1)$ is $\mathbf{DM}_{\text{gm}}(S)$. In other words, $\mathbf{DM}_{\text{gm}}(S)$ is obtained from $\mathbf{DM}_{\text{gm}}^{\text{eff}}(S)$ by tensor-inverting $\mathbf{1}(-1)$.

Definition 1.7. Let M be any geometric motive over S . We write $H^i(M) := H^i(S, M) := \text{Hom}_{\mathbf{DM}(S)}(\mathbf{1}, M[i])$. For $M = M(X)(n)$ for any X over S we also write $H^i(X, n) := H^i(M(X)(n)) = \text{Hom}_{\mathbf{DM}_{\text{gm}}(S)}(\mathbf{1}, M(X)(n)[i])$. This is called *motivic cohomology* of M and X , respectively.

Axiom 1.8. (Motivic cohomology vs. K -theory) For any regular scheme X , there is an isomorphism $H^i(X, n) \cong K_{2n-i}(X)_{\mathbb{Q}}^{(n)}$, where the right hand term denotes the Adams eigenspace of algebraic K -theory tensored with $\mathbb{Q}$ [Qui73].

This is a key property of motives, since algebraic K -theory is a universal cohomology theory in the sense that Chern characters map from algebraic K -theory to any other (reasonable) cohomology theory of algebraic varieties [Gil80]. For S a perfect field, this axiom is given by [Voe00, Prop. 4.2.9] and its non-effective analogue. See also [Lev98, Theorem I.III.3.6.12.].

Recall Grothendieck's category of pure motives $\mathbf{M}_{\sim}(K)$ with respect to an adequate equivalence relation $\sim$, see e.g. [And04, Section 4]. For rational equivalence they are also called Chow motives, since, for any smooth projective variety X over a field K ,

$$\mathrm{Hom}_{\mathbf{M}_{\mathrm{rat}}(K)}(\mathbf{1}(-n), h(X)) = \mathrm{CH}^n(X), \quad (1)$$

where $h(X)$ denotes the Chow motive of X and the right hand term is the *Chow group* of cycles of codimension n in X . This way, the above axiom models the fact [Voe00, 2.1.4] that Chow motives are a full subcategory of $\mathbf{DM}_{\mathrm{gm}}(K)$. Under the embedding $\mathbf{M}_{\mathrm{rat}}(K) \subset \mathbf{DM}_{\mathrm{gm}}(K)$, $h(X, n)$ maps to $\mathrm{M}(X)(n)[2n]$.

Remark 1.9. We do not need to assume *expressis verbis* homotopy invariance (i.e., $\mathbf{1} \xrightarrow{\cong} pr_* pr^* \mathbf{1} \in \mathbf{DM}_{\mathrm{gm}}(S)$ for $pr : S \times \mathbb{A}^1 \rightarrow S$) nor the projective bundle formula [Voe00, Prop. 3.5.1]. (Note, however, that K' -theory does have such properties.)

Axiom 1.10. (Localization) Let $i : Z \rightarrow S$ be any closed immersion and $j : V \rightarrow S$ the open complement. The adjointness maps give rise to the following distinguished triangles in $\mathbf{DM}(S)$:

$$\begin{aligned} j_! j^* &\rightarrow \mathrm{id} \rightarrow i_* i^*, \\ i_* i^! &\rightarrow \mathrm{id} \rightarrow j_* j^*. \end{aligned}$$

(In particular, $f_* f^* \cong \mathrm{id}$, where $f : X_{\mathrm{red}} \rightarrow X$ denotes the canonical map of the reduced subscheme structure.) In addition, one has $j^* j_* = \mathrm{id}$ and $i^* i_* = \mathrm{id}$, equivalently $j^* i_* = i^* j_! = 0$.

It should be emphasized that the localization axiom requires rational coefficients.

Axiom 1.11. (Purity and base change)

- For any quasi-projective map f , there is a functorial transformation of functors $f_! \rightarrow f_*$. It is an isomorphism if f is projective.
- (*Relative purity*): If f is quasi-projective and smooth of constant relative dimension d , there is a functorial (in f) isomorphism $f^! \cong f^*(d)[2d]$.

- (*Absolute purity*): If $i : Z \rightarrow U$ is a closed immersion of codimension c of two regular schemes Z and U , there is a natural isomorphism $i^! \mathbf{1} \cong \mathbf{1}(-c)[-2c]$.
- (*Base change*): For any two quasi-projective maps f and g let f' and g' denote the pullback maps:

$$\begin{array}{ccc} X' \times_X Y & \xrightarrow{g'} & Y \\ \downarrow f' & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array} \quad (2)$$

Then there is canonical isomorphism of functors

$$f^* g! \xrightarrow{\cong} g'_! f'^*.$$

This axiom is proven by Cisinski & Déglise using Ayoub's general base change formalism. See in particular [Ayo07, 1.4.11, 12] for the construction of the base change map. See also [Lev98, Theorem I.I.2.4.9] for a similar statement in Levine's category of motives.

Definition 1.12. Let $f : S \rightarrow \text{Spec } \mathbb{Z}$ be the structural map. Assume f is quasi-projective. Then $D(M) := \underline{\text{Hom}}(M, f^! \mathbf{1}(1)[2])$ is called *Verdier dual* of M .

By the preceding axioms, D induces a contravariant endofunctor of $\mathbf{DM}_{\text{gm}}(S)$. The shift and twist in the definition is motivated as follows: given some complex analytic space X , the Verdier dual of a sheaf $\mathcal{F}$ on X is defined by

$$D(\mathcal{F}) := \underline{\text{RHom}}_{\mathbf{D}(\text{Shv}(X))}(\mathcal{F}, f^! \mathbb{Z}),$$

where f denotes the projection to a point, see e.g. [Ive86, Ch. VI]. When X is smooth of dimension d , one has $f^! \mathbb{Z} = f^* \mathbb{Z}(d)[2d] = \mathbb{Z}(d)[2d]$. A similar fact holds for ℓ -adic sheaves (see e.g. [KW01, Section II.7-8]). The above definition mimics this situation insofar as $\text{Spec } \mathbb{Z}$ is seen as an analogue of a smooth affine curve.

Let us give a number of consequences of the preceding axioms, in particular purity, base change and localization: in (2), suppose that f is smooth and $X' \subset X$ is a codimension one closed immersion between regular schemes. Then there is a canonical isomorphism

$$i^! M_S(X) = M_Z(X \times_S Z)(-1)[-2]. \quad (3)$$

Let $Z \subset X$ be a closed immersion of quasiprojective schemes over S . Then there is a distinguished triangle of motives with compact support

$$M_c(Z) \rightarrow M_c(X) \rightarrow M_c(X \setminus Z).$$

Let S be a regular scheme of equidimension d that is affine or projective over $\text{Spec } \mathbb{Z}$. Let $f : S \rightarrow \text{Spec } \mathbb{Z}$ be the structural map. Then $f^! \mathbf{1} = \mathbf{1}(d-1)[2d-2]$, as one sees by applying relative and absolute purity to the maps p and i , where $f = p \circ i$ is a factorization into a affine or projective projection map and a closed immersion. In particular, the Verdier duality functor on any open subscheme S of $\text{Spec } \mathcal{O}_F$ is given by $D_{\mathbf{DM}_{\text{gm}}(S)}(?) = \underline{\text{Hom}}(?, \mathbf{1}(1)[2])$ while on $\mathbf{DM}_{\text{gm}}(\mathbb{F}_{\mathfrak{p}})$ it is given by $\underline{\text{Hom}}(?, \mathbf{1}) = ?^{\vee}$.

Axiom 1.13. (Verdier dual) The Verdier dual functor D exchanges “ $!$ ” and “ $*$ ” throughout, e.g., there are natural isomorphisms $D(f^! M) \cong f^* D(M)$ for any quasi-projective map $f : X \rightarrow Y$ and $M \in \mathbf{DM}(Y)$ and similarly with $f_!$ and f_* .

Lemma 1.14. *Let S be such that $f^! \mathbf{1} = f^* \mathbf{1}(d)[2d]$ for some integer d , where $f : S \rightarrow \text{Spec } \mathbb{Z}$ is the structural map. For example, S might be regular and affine or projective over $\mathbb{Z}$ (see above), or smooth over $\text{Spec } \mathbb{Z}$ (purity). Then, for any compact object $M \in \mathbf{DM}_{\text{gm}}(S)$, the canonical map $M \rightarrow D(D(M))$ is an isomorphism. This will be referred to as reflexivity of Verdier duality.*

Proof: By Axiom 1.5, it suffices to check it for $M = \pi_* \pi^* \mathbf{1}$, where $\pi : X \rightarrow S$ is some map of finite type. In this case it follows for adjointness reasons and the assumption. $\square$

Axiom 1.15. (Resolution of singularities) Let K be a field. As a triangulated additive tensor category (i.e., closed under triangles, arbitrary direct sums and tensor product), $\mathbf{DM}(K)$ is generated by $\mathbf{1}(-1)$ and all $M(X)$, where X/K is a smooth projective variety.

When $S = \text{Spec } \mathcal{O}_F$, the generators of $\mathbf{DM}(S)$ are $\mathbf{1}(-1)$, $i_{\mathfrak{p}*} M(X_{\mathfrak{p}})$, and $M(X)$, instead, where $X_{\mathfrak{p}}$ is any projective and smooth variety over $\mathbb{F}_{\mathfrak{p}}$, $i_{\mathfrak{p}}$ denotes the immersion of any closed point $\mathbb{F}_{\mathfrak{p}}$ of S , and X is any regular, flat projective scheme over $\mathcal{O}_F$.

Consequently, the subcategories of compact objects $\mathbf{DM}_{\text{gm}}(-)$ are generated as a thick tensor subcategory by the mentioned objects. In Voevodsky’s theory of motives over a field of characteristic zero, this is [Voe00, Section 4.1]. This uses Hironaka’s resolution of singularities. Over a field of positive characteristic and number rings, one has to use de Jong’s resolution result, see [HK06, Lemma B.4].

We also need a limit property of the generic point. Let S be an open subscheme of $\text{Spec } \mathcal{O}_F$, let $\eta : \text{Spec } F \rightarrow S$ be the generic point.

Axiom 1.16. (Generic point) Let M be any geometric motive over S . The natural maps $j_* j^* M \rightarrow \eta_* \eta^* M$ give rise to an isomorphism $\varinjlim j_* j^* M = \eta_* \eta^* M$, where the colimit is over all open subschemes $j : S' \rightarrow S$. It induces a distinguished triangle in $\mathbf{DM}(S)$

$$\bigoplus_{\mathfrak{p} \in S} i_{\mathfrak{p}*} i_{\mathfrak{p}}^! M \rightarrow \text{id} \rightarrow \eta_* \eta^* M, \quad (4)$$

where the sum runs over all closed points $\mathfrak{p} \in S$ and $i_{\mathfrak{p}}$ is the closed immersion.

2 Realizations

One of the main interests in motives lies in the fact that they are explaining (or are supposed to explain) common phenomena in various cohomology theories. These cohomology functors are commonly referred to as *realization* functors. They typically have the form $\mathbf{DM}_{\text{gm}}(S) \rightarrow \mathbf{D}^b(\mathcal{C})$, where $\mathcal{C}$ is an abelian category whose objects are amenable with the methods of (linear) algebra, such as finite-dimensional vector spaces or finite-dimensional continuous group representations or constructible sheaves.

For example, let ℓ be a prime and let S be either a field of characteristic different from ℓ or a scheme of finite type over $\text{Spec } \mathcal{O}_F[1/\ell]$. The ℓ -adic cohomology maps any scheme X over S to

$$\mathbf{R}\Gamma_\ell(X) := \mathbf{R}\pi_* \pi^* \mathbb{Q}_\ell \in \mathbf{D}_c^b(S, \mathbb{Q}_\ell),$$

where $\pi : X \rightarrow S$ is the structural map and the right hand category denotes the “derived” category of constructible $\mathbb{Q}_\ell$ -sheaves on S (committing the standard abuse of notation, see e.g. [KW01, II.6., II.7.]). This functor factors over the ℓ -adic realization functor ([Hub00, p. 772], [Ivo07]) $\mathbf{R}\Gamma_\ell : \mathbf{DM}_{\text{gm}}(S) \rightarrow \mathbf{D}_c^b(S, \mathbb{Q}_\ell)$. When S is of finite type over $\mathbb{F}_p$, the realization functor actually maps to $\mathbf{D}_{c,m}^b(S, \mathbb{Q}_\ell)$, the full subcategory of complexes C in $\mathbf{D}_c^b(S, \mathbb{Q}_\ell)$ such that all $H^n(C)$ are mixed sheaves [Del80, 1.2].

Further realization functors include Betti, de Rham and Hodge realization. See e.g. [Hub00, 2.3.5]. The following axiom says (in particular) that the ℓ -adic realization of $M(X)$ does give the ℓ -adic cohomology groups.

Axiom 2.1. (Functoriality and realizations) The ℓ -adic realization functor commutes with the six Grothendieck functors f_* , $f_!$, $f^!$, f^* , $\otimes$ and $\underline{\text{Hom}}$ (where applicable). For example, for any map $f : S' \rightarrow S$ and any geometric motive M over S' :

$$(f_* M)_\ell = f_*(M_\ell).$$

3 Interlude: Perverse sheaves over number rings

This section is devoted to a modest extension of ℓ -adic perverse sheaves [BBD82] to the situation where the base S is an open subscheme of $\text{Spec } \mathcal{O}_F[1/\ell]$. It is needed to formulate Axiom 4.8 for the ℓ -adic realization of motives over number rings. In a nutshell, the theory of perverse sheaves on varieties over $\mathbb{F}_q$ stakes on relative purity, that is $f^! \mathbb{Z}_\ell = f^* \mathbb{Z}_\ell(n)[2n]$ for a smooth map f of relative dimension n . The analogous identity for a closed immersion $i : \text{Spec } \mathbb{F}_p \rightarrow S$ reads

$$i^! \mathbb{Z}_\ell = i^* \mathbb{Z}_\ell(-1)[-2]. \quad (5)$$

It is a reformulation of well-known cohomological properties of the inertia group: $H^1(I_p, V) = (V(-1))_{I_p}$ for any ℓ -adic module with continuous I_p -action ($p \nmid \ell$). All higher group cohomologies of I_p vanish.

Let $\mathbf{D}^b(S, \mathbb{Z}_\ell)$ be the bounded “derived” category of $\mathbb{Z}_\ell$ -sheaves on S as constructed by Ekedahl [Eke90]. All following constructions can be done for $\mathbb{Q}_\ell$ instead of $\mathbb{Z}_\ell$, as well. We keep writing j_* for the total derived functor, commonly denoted Rj_* etc. However, $R^n j_*$ etc. keep their original meaning.

As in *loc. cit.*, see especially [2.2.10, 2.1.2, 2.1.3, 1.4.10]¹, one first defines a notion of stratification, and secondly obtains a t -structure on the subcategory $\mathbf{D}_{(\Sigma, L)}^b(S, \mathbb{Z}_\ell)$ that are constructible with respect to a given stratification $\Sigma = \{\Sigma_i\}$ and a set L of irreducible lisse sheaves on the strata. Thirdly, one takes the “limit” over the stratifications. The union of all $\mathbf{D}_{(\Sigma, L)}^b(S, \mathbb{Z}_\ell)$ is the “derived” category $\mathbf{D}_c^b(S, \mathbb{Z}_\ell)$ of constructible sheaves. In order to extend the t -structure on the subcategories to one on $\mathbf{D}_c^b(S, \mathbb{Z}_\ell)$, one has to check that the inclusion $\mathbf{D}_{(\Sigma', L')}^b(S, \mathbb{Z}_\ell) \rightarrow \mathbf{D}_{(\Sigma, L)}^b(S, \mathbb{Z}_\ell)$ is t -exact for any refinement of stratifications. Here we employ a different argument. The proof of [2.1.14, 2.2.11] relies on relative purity for ℓ -adic sheaves [SGA73, Exp. XVI, 3.7]. As in the proof of [2.1.14] we have to check the following: let $\Sigma_i \xrightarrow{a} \Sigma'_i \xrightarrow{b} S$ be the inclusions of some strata and let $C \in \mathbf{D}_{(\Sigma', L')}^{b, \geq 0}(S, \mathbb{Z}_\ell)$. Then $C \in \mathbf{D}_{(\Sigma, L)}^{b, \geq 0}(S, \mathbb{Z}_\ell)$. We can assume $\dim \Sigma_i = 0$, $\dim \Sigma'_i = 1$, since all other cases are clear. Thus, b is an open immersion. We may also assume for notational simplicity that $\Sigma_i = \text{Spec } \mathbb{F}_p$. Let j be the complementary open immersion to a . By definition, $H^n b^! C = b^! H^n C = b^* H^n C$ is locally constant and vanishes for $n < -1$. In the parlance of Galois modules this means that, viewed as a $\pi_1(\Sigma'_i)$ -representation, the action of the inertia group $I_p \subset \pi_1(\Sigma'_i)$ on that sheaf is trivial. Thus

$$a^! H^n b^* C = a^*(R^1 j_* j^* H^n b^* C)[-2] = H^1(I_p, H^n b^* C)[-2] = a^* H^n b^* C(-1)[-2].$$

(We have used $p \nmid \ell$ at this point.) The spectral sequence

$$H^{p-2} a^* H^q b^! C(-1) = H^p a^! H^q b^! C \Rightarrow H^n a^! b^! C$$

is such that the left hand term vanishes for $p \neq 2$ since a^* is exact w.r.t. the standard t -structure. It also vanishes for $q < -1$ by the above. Hence the right hand term vanishes for $n = p+q < 1$. A fortiori it vanishes for $n < -\dim \mathbb{F}_p = 0$.

Objects in the heart of this t -structure on $\mathbf{D}_c^b(S, \mathbb{Z}_\ell)$ are called *perverse sheaves* on S . For example $\mathbb{Z}_\ell[1]$ and $i_* \mathbb{Z}_\ell$ for any immersion i of a closed point are perverse sheaves on S . The *Verdier dual* of any $C \in \mathbf{D}_c^b(S, \mathbb{Z}_\ell)$ is defined by $D(C) := \underline{\text{Hom}}(C, \mathbb{Z}_\ell(1)[2])$. As above, we have dropped “ R ” from the notation, so that this Hom means what is usually denoted RHom.

Lemma 3.1. *Let $j : S' \rightarrow S$ be an open immersion and $i : Z \rightarrow S$ a closed immersion. Let $\eta : \text{Spec } F \rightarrow S$ be the generic point. Then j_* , $j_!$, i_* , $\eta^*[-1]$, j^* and D are t -exact, while i^* ($i^!$) is of cohomological amplitude $[-1, 0]$ ($[0, 1]$), in particular right-exact (left-exact, respectively). Finally, the t -structure on $\mathbf{D}_c^b(S, \mathbb{Z}_\ell)$ is non-degenerate.*

Proof: The only non-formal statement is the exactness of j_* . The corresponding precursor result [4.1.10] is a reformulation of [SGA72, Th. 3.1., Exp.

¹In the sequel, any reference in brackets refers to [BBD82].

XIV], which says for any affine map $j : X \rightarrow Y$ over schemes over a field K , and any (honest) sheaf $\mathcal{F}$ which is torsion (prime to $\text{char } K$)

$$d(\mathbf{R}^q j_* \mathcal{F}) \leq d(\mathcal{F}) - q$$

where $d(\mathcal{G}) := \sup\{\dim \overline{\{x\}}, \mathcal{G}_{\overline{x}} \neq 0\}$ for any sheaf $\mathcal{G}$. In our situation, we are given a locally constant sheaf $\mathcal{F}$ on S' whose torsion is prime to all characteristics of S . The conclusion of the theorem also holds for j , as follows from the cohomological dimension of $I_{\mathbf{p}}$, which is one. $\square$

Let $\mathcal{F}$ be any perverse sheaf on S' . Following [1.4.22], let the *intermediate extension* $j_{!*} \mathcal{F}$ be the image of the map $j_! \mathcal{F} \rightarrow j_* \mathcal{F}$ of perverse sheaves on S . As in [2.1.11] one sees that it can be calculated in terms of the good truncation with respect to the standard t -structure: $j_{!*} \mathcal{F} = \tau_{\leq -1}^{can} j_* \mathcal{F}$. If $\mathcal{F} = \mathcal{G}[1]$, where $\mathcal{G}$ is a lisse (honest) sheaf on S' , this gives $(\mathbf{R}^0 j_* \mathcal{G})[1]$.

4 Mixed motives

Throughout this section, let $S = \text{Spec } F$ or $\text{Spec } \mathbb{F}_{\mathbf{p}}$ or an open subscheme of $\text{Spec } \mathcal{O}_F$.

This section formulates a number of axioms concerning weights and the motivic t -structure on triangulated categories of motives over S . In contrast to the axioms listed in Section 1, the axioms mentioned in this section are wide open, so it might be more appropriate to call them conjectures instead.

Axiom 4.1. (Motivic t -structure and cohomological dimension) The category of geometric motives $\mathbf{DM}_{\text{gm}}(S)$ has a non-degenerate t -structure [BBD82, Def. 1.3.1] called *motivic t -structure*. Its heart is denoted $\mathbf{MM}(S)$. Objects of $\mathbf{MM}(S)$ are called *mixed motives* over S .

For any $M \in \mathbf{DM}_{\text{gm}}(S)$, there are $a, b \in \mathbb{Z}$ such that $\tau_{\leq a} M = \tau_{\geq b} M = 0$. Here and in the sequel, $\tau_{\leq -}$ and $\tau_{\geq -}$ denote the truncation functors with respect to the motivic t -structure.

The *cohomological dimension* of $\mathbf{DM}_{\text{gm}}(\mathbb{F}_{\mathbf{p}})$ and $\mathbf{DM}_{\text{gm}}(F)$ is 0 and 1, respectively, in the sense that

$$\text{Hom}_{\mathbf{DM}(\mathbb{F}_{\mathbf{p}})}(M, N[i]) = 0$$

for all mixed motives M, N over $\mathbb{F}_{\mathbf{p}}$ and $i > 0$ and similarly for mixed motives over F and $i > 1$. (For $i < 0$ the term vanishes by the t -structure axioms.)

The t -structures are such that over $S = \text{Spec } F$ or $\text{Spec } \mathbb{F}_{\mathbf{p}}$, $\mathbf{1} \in \mathbf{MM}(S)$, while for an open subscheme $S \subset \text{Spec } \mathcal{O}_F$, $\mathbf{1}[1] \in \mathbf{MM}(S)$.

The existence of the motivic t -structure on $\mathbf{DM}_{\text{gm}}(K)$ satisfying the axioms listed in this section is part of the general motivic conjectural framework, see e.g. [Beĭ87a, App. A], [And04, Ch. 21]. The idea of building a triangulated category of motives and descending to mixed motives by means of a t -structure

is due to Deligne. The existence of a motivic t -structure on $\mathbf{DM}_{\text{gm}}(K)$ is only known in low dimensions: the subcategory of Artin motives, i.e., motives of zero-dimensional varieties, carries such a t -structure [Voe00, Section 3.4.]. By *loc. cit.*, [Org04], the subcategory of $\mathbf{DM}_{\text{gm}}(K)$ generated by motives of smooth varieties of dimension ≤ 1 is equivalent to the bounded derived category of 1-motives [Del74, Section 10] up to isogeny. Finally, if K is a field satisfying the Beilinson-Soulé vanishing conjecture, such as a finite field or a number field, the category of Artin-Tate motives over K enjoys a motivic t -structure [Lev93, Wil]. The results on Artin-Tate motives are generalized to bases S which are open subschemes of $\text{Spec } \mathcal{O}_F$ in [Scha].

The conjecture about the cohomological dimension is due to Beilinson. A (fairly weak) evidence for this conjecture is the cohomological dimension of Tate motives over F and $\mathbb{F}_p$, which is one and zero, respectively. This follows from vanishing properties of K -theory of these fields.

The normalization in the last item is merely a matter of bookkeeping, but is motivated by similar shifts in perverse sheaves (Section 3). The existence of a motivic t -structure is not expected to hold for motives with integral coefficients.

We do not (need to) assume that the canonical functor $\mathbf{D}^b(\mathbf{MM}(S)) \rightarrow \mathbf{DM}_{\text{gm}}(S)$ is an equivalence of categories or, equivalently [Beĭ87b, Lemma 1.4.], $\text{Ext}_{\mathbf{MM}(S)}^i(A, B) = \text{Hom}_{\mathbf{DM}_{\text{gm}}(S)}(A, B[i])$ for all mixed motives A and B .

Axiom 4.2. (Exactness properties) Let $S \subset \text{Spec } \mathcal{O}_F$ be an open subscheme, let $i : \text{Spec } \mathbb{F}_p \rightarrow \text{Spec } \mathcal{O}_F$ be a closed point, $j : U \rightarrow S$ an open immersion and $\eta : \text{Spec } F \rightarrow S$ the generic point.

Then $j^* = j^!$, $\eta^*[-1]$, i_* , j_* and $j_!$ are exact with respect to the motivic t -structures on the involved categories of geometric motives. Further, i^* is right-exact, more precisely it maps objects in cohomological degree 0 to degrees $[-1, 0]$. Dually, $i^!$ has cohomological amplitude $[0, 1]$. Verdier duality D is “anti-exact”, i.e., maps objects in positive degrees to ones in negative degrees and vice versa.

The axiom is motivated by the same exactness properties in the situation of perverse sheaves over $\text{Spec } \mathcal{O}_F[1/\ell]$ (Section 3). The corresponding exactness properties of the above functors on Artin-Tate motives, where the motivic t -structure is available, is established in [Scha].

Definition 4.3. The cohomology functor with respect to the motivic t -structure on $\mathbf{DM}_{\text{gm}}(S)$ is denoted ${}^p\mathbf{H}^*$. For any scheme X/S , we write

$$\mathbf{h}^i(X, n) := {}^p\mathbf{H}^i \mathbf{M}_S(X)(n).$$

Axiom 4.4. (Hard Lefschetz) Let X_η/F be smooth and projective of constant dimension d_η . Let $i \leq d_\eta$ and a any integer. Then, taking the $(d_\eta - i)$ -fold cup product with the cycle class of a hyperplane section with respect to an embedding of X_η into some projective space over F yields an isomorphism (“hard Lefschetz isomorphism”)

$$\mathbf{h}^i(X_\eta, a) \xrightarrow{\cong} \mathbf{h}^{2d_\eta-i}(X_\eta, d_\eta - i + a).$$

Axiom 4.5. (Decomposition of smooth projective varieties) Let X/S be smooth and projective. In $\mathbf{DM}_{\text{gm}}(S)$, there is a non-canonical isomorphism

$$\phi_X : M(X) \cong \bigoplus_n h^n(X)[-n].$$

For open subschemes $S \subset \text{Spec } \mathcal{O}_F$, this isomorphism is compatible with pull-backs along all closed points $i : \text{Spec } \mathbb{F}_{\mathfrak{p}} \rightarrow S$ in the following sense: let $X_{\mathfrak{p}}$ be the fiber of X over $\mathbb{F}_{\mathfrak{p}}$, and let ψ be the isomorphism making the following diagram commutative. Its left hand isomorphism is an instance of base change.

$$\begin{array}{ccc} i^* M(X) & \xrightarrow{i^* \phi_X} & \bigoplus_n i^* h^n(X)[-n] \\ \downarrow \cong & & \downarrow \psi \\ M(X_{\mathfrak{p}}) & \xrightarrow{\phi_{X_{\mathfrak{p}}}} & \bigoplus_m h^m(X_{\mathfrak{p}})[-m] \end{array}$$

Then ψ respects the direct summands, i.e., induces isomorphisms

$$i^* h^n(X)[-n] \cong h^{n-1}(X_{\mathfrak{p}})[-n+1].$$

Remark 4.6. Axiom 4.4 is part of the classical conjectural framework of motives (see e.g. [And04, Section 5.2, Prop. 5.2.5.1]). It implies the decomposition of Axiom 4.5 [Del94]. The latter is used in Lemma 5.11, which in turn is crucial in Section 6. The index shift in Axiom 4.5 is due to the normalization in Axiom 4.1: for $S = \text{Spec } \mathcal{O}_F$ and a closed point i as above, take for example $X = S$, $M(S) = \mathbf{1} = h^1(S)[-1]$ (sic) and $i^* M(S) = \mathbf{1}_{\mathbb{F}_{\mathfrak{p}}} = h^0(\text{Spec } \mathbb{F}_{\mathfrak{p}})$.

Consider any smooth projective variety X_{η} over F . Using that the cohomological dimension of $\mathbf{DM}_{\text{gm}}(F)$ is one, the term $H^{2n}(X_{\eta}, n)$, that is identified with $\text{CH}^n(X_{\eta})_{\mathbb{Q}}$ by Axiom 1.8, appears in an exact sequence of the following form

$$0 \rightarrow H^1(h^{2n-1}(X_{\eta}, n)) \rightarrow H^{2n}(X_{\eta}, n) \rightarrow H^0(h^{2n}(X_{\eta}, n)) \rightarrow 0.$$

Consider the cycle class map from the m -th Chow group to ℓ -adic cohomology of X_{η} , $\text{CH}^m(X_{\eta})_{\mathbb{Q}} \rightarrow H^{2m}(X_{\eta}, \mathbb{Q}_{\ell}(m))$ [Mil80, VI.9]. The groups $\text{CH}^m(X_{\eta})_{\mathbb{Q}, \text{hom}}$ and $\text{CH}^m(X_{\eta})_{\mathbb{Q}}/\text{hom}$ are by definition the kernel and the image (seen as a quotient of the Chow group) of this map. According to standard conjectures, homological equivalence should not depend on the choice of the cohomology theory, since they are all supposed to agree with numerical equivalence, see e.g. [And04, Chapter 5].

Axiom 4.7. (Morphisms in $\mathbf{MM}(F)$ and Chow groups) For such X_{η} , the sequence above identifies with

$$0 \rightarrow \text{CH}^n(X_{\eta})_{\mathbb{Q}, \text{hom}} \rightarrow \text{CH}^n(X_{\eta})_{\mathbb{Q}} \rightarrow \text{CH}^n(X_{\eta})_{\mathbb{Q}}/\text{hom} \rightarrow 0.$$

Let either S be a field and let C stand for the ℓ -adic realization (in case $\text{char } S \neq \ell$), Betti, de Rham or absolute Hodge realization or let $S \subset \text{Spec } \mathcal{O}_F[1/\ell]$ be an open subscheme and let C be the ℓ -adic realization. We write $\text{R}\Gamma_C : \mathbf{DM}_{\text{gm}}(S) \rightarrow \mathbf{D}^b(C)$ for the realization functor, where $\mathbf{D}^b(C)$ is understood as a placeholder of the target category of C . (We abuse the notation insofar as that target category is not a derived category in the strict sense when C is the ℓ -adic realization.) For all realizations over a field, this category is endowed with the usual t -structure, e.g. on $\mathbf{D}_c^b(K, \mathbb{Q}_\ell)$ for ℓ -adic realization. When S is an open subscheme of $\text{Spec } \mathcal{O}_F[1/\ell]$ and C is the ℓ -adic realization, we take the perverse t -structure on $\mathbf{D}_c^b(S, \mathbb{Q}_\ell)$ defined in Section 3. Using this, we have the following axiom:

Axiom 4.8. (Exactness of realization functors) Realization functors $\text{R}\Gamma_C$ are exact with respect to the motivic t -structure on $\mathbf{DM}_{\text{gm}}(S)$. Equivalently, as the t -structure on $\mathbf{D}^b(C)$ is non-degenerate, $\text{R}\Gamma_C({}^p\text{H}^0 M) = {}^p\text{H}^0 \text{R}\Gamma_C(M)$ for any geometric motive M over S . On the left, ${}^p\text{H}^0$ denotes the cohomology functor belonging to the motivic t -structure on $\mathbf{DM}_{\text{gm}}(S)$, while on the right hand side it is the one belonging to the t -structure on $\mathbf{D}^b(C)$.

This axiom is, if fairly loosely, motivated by a similar fact in the theory of mixed Hodge modules: let X be any complex algebraic variety. Then, under the faithful “forgetful functor” from the derived category of mixed Hodge modules to the derived category of constructible sheaves with rational coefficients

$$\mathbf{D}^b(\mathbf{MHM}(X)) \rightarrow \mathbf{D}_c^b(X, \mathbb{Q})$$

the category $\mathbf{MHM}(X)$ corresponds to perverse sheaves on X .

Recall that in an abelian category $\mathcal{C}$, a morphism $f : (X, W^*) \rightarrow (Y, W^*)$ between filtered objects is called *strict* if $f(W^n X) = f(X) \cap W^n Y$ for all n .

Axiom 4.9. (Weights) Any mixed motive M over S has a functorial finite exhaustive separated filtration $W_* M$ called *weight filtration*, i.e., a sequence of subobjects in the abelian category $\mathbf{MM}(S)$

$$0 = W_a M \subset W_{a+1} M \subset \cdots \subset W_b M = M.$$

Any morphism between mixed motives is strict with respect to the weight filtration.

Given two motives of pure weight, their tensor product is pure; its weight is the sum of the two individual weights. In particular, the unit motive $\mathbf{1}$ is pure of weight zero.

Let $\text{R}\Gamma_C : \mathbf{DM}_{\text{gm}}(S) \rightarrow \mathbf{D}^b(C)$ be any realization functor that has a notion of weights (such as the ℓ -adic realization when $S = \text{Spec } \mathbb{F}_p$ or the Hodge realization when $S = \text{Spec } \mathbb{Q}$). Then

$$\text{gr}_n^W \text{R}\Gamma_C(M) = \text{R}\Gamma_C(\text{gr}_n^W M)$$

for any mixed motive M over S .

Definition 4.10. For any $M \in \mathbf{MM}(S)$, we write $\mathrm{wt}(M)$ for the (finite) set of integers n such that $\mathrm{gr}_n^W M \neq 0$. For $M \in \mathbf{DM}_{\mathrm{gm}}(S)$, define $\mathrm{wt}(M) := \bigcup_{i \in \mathbb{Z}} \mathrm{wt}(\mathrm{PH}^i(M)) - i$.

Axiom 4.11. (Preservation of weights) Let $f : X \rightarrow S$ be a quasi-projective map. Then the functors $f_! f^*$ preserve negativity of weights, i.e., given a geometric motive M over S with weights ≤ 0 , $f_! f^* M$ also has weights ≤ 0 . Dually, $f_* f^!$ preserves positive weights.

In the particular case $S \subset \mathrm{Spec} \mathcal{O}_F$ (open), let $j : U \rightarrow S$ and $\eta : \mathrm{Spec} F \rightarrow S$ be an open immersion into S and the generic point of S , respectively. Let $i : \mathrm{Spec} \mathbb{F}_p \rightarrow S$ be a closed point. Then, i^* and $j_!$ preserve negativity of weights and dually for $i^!$ and j_* . Finally, j^* and η^* both preserve both positivity and negativity of weights.

The preceding weight axioms are motivated by the very same properties of ℓ -adic perverse sheaves on schemes over $\mathbb{C}$ or finite fields [BBD82, 5.1.14], number fields [Hub97] as well as Hodge structures [Del74, Th. 8.2.4] and Hodge modules (see [PS08, Chapter 14.1] for a synopsis). In these settings, actually $f_!$ and f^* preserve negative weights, but we do not need weights for motives over more general bases than the ones above. The weight formalism we require is stronger than the one provided by the differential-graded interpretation of $\mathbf{DM}_{\mathrm{gm}}$ over a field [Bon09] or [BV08, 6.7.4].

Example 4.12. For any projective (smooth) scheme X of finite type over S , the weights of $\mathrm{h}^i(X)(n)$ are $\leq i - 2n$ ($\geq i - 2n$, respectively).

Axiom 4.13. (Mixed vs. pure motives) For any field K , the subcategory of pure objects in $\mathbf{MM}(K)$ identifies with $\mathbf{M}_{\mathrm{num}}(K)$, the category of numerical pure motives over K .

As a consequence of the weight filtration, every mixed motive is obtained in finitely many steps by taking extensions of motives in $\mathbf{M}_{\mathrm{num}}(K)$.

5 Motives over number rings

In the following sections we assume the axioms of Sections 1, 2, and 4. Unless explicitly mentioned otherwise, let S be an open subscheme of $\mathrm{Spec} \mathcal{O}_F$, let $i : \mathrm{Spec} \mathbb{F}_p \rightarrow \mathrm{Spec} \mathcal{O}_F$ be a closed point, $j : S' \rightarrow S$ an open subscheme and $\eta : \mathrm{Spec} F \rightarrow S$ the generic point.

This section derives a number of basic results about motives over S from the axioms spelled out above. We define and study the *intermediate extension* $j_{!*} : \mathbf{MM}(S') \rightarrow \mathbf{MM}(S)$ in analogy to perverse sheaves (Definition 5.3). An “explicit” set of generators of $\mathbf{DM}_{\mathrm{gm}}(S)$ (Proposition 5.7) is obtained using $j_{!*}$. We introduce a notion of *smooth motives* (Definition 5.8), which should be thought of as analogs of lisse sheaves. Using this notion, we extend the intermediate extension to a functor $\eta_{!*}$ spreading out motives over F with a certain smoothness property to motives over S , cf. Definition 5.14. This functor

will be the main technical tool in dealing with f -cohomology in Section 6. In Lemma 5.16 we express the ℓ -adic realization of motives of the form $j_{!*}M$ in sheaf-theoretic language.

5.1 Cohomological dimension

The following is an immediate consequence of Axiom 4.2:

Lemma 5.1. *For any scheme X over S we have $\eta^*[-1]h^i(X, n) = h^{i-1}(X \times_S F, n)$.*

The following lemma parallels (and is a consequence of) Axiom 4.1.

Lemma 5.2. *The cohomological dimension of $\mathbf{DM}_{\text{gm}}(S)$ is two, that is to say, for any two mixed motives M, N over S ,*

$$\text{Hom}_{\mathbf{DM}_{\text{gm}}(S)}(M, N[i]) = 0$$

for all $i > 2$. In particular $H^i(M)$ vanishes for $|i| > 1$.

Proof: Apply $\text{Hom}(M, -)$ to the localization triangle $\bigoplus_{p \in S} i_{p*} i_p^! N \rightarrow N \rightarrow \eta_* \eta^* N$ of Axiom 1.16, where i_p are the immersions of the closed points of S . The terms adjacent to $\text{Hom}(M, N[i])$ are $\text{Hom}(M, \bigoplus_p i_{p*} i_p^! N[i]) = \bigoplus_p \text{Hom}(i_p^* M, i_p^! N[i])$ (as M is compact) and $\text{Hom}(M, \eta_* \eta^* N[i]) = \text{Hom}(\eta^* M, \eta^* N[i])$. The latter term vanishes for $i > 1$ since $\eta^*[-1]$ is exact and the cohomological dimension of $\mathbf{DM}_{\text{gm}}(F)$ is one.

To deal with the former term, we have to take into account that $i_p^!$ and i_p^* are not t -exact, but of cohomological amplitude $[0, 1]$ and $[-1, 0]$, respectively. By decomposing $i_p^! N$ into its ${}^p\text{H}^{-1}$ - and ${}^p\text{H}^0$ -part and similarly with $i_p^* M$ and using that the cohomological dimension of $\mathbf{DM}_{\text{gm}}(\mathbb{F}_p)$ is zero, the term vanishes for $i > 2$. Using general t -structure properties, the second claim is a particular case of the first one. $\square$

5.2 Intermediate extension

Definition 5.3. (Motivic analog of [BBD82, Def. 1.4.22]) The *intermediate extension* $j_{!*}$ of some mixed motive M over S' is defined as

$$j_{!*}M := \text{im}(j_!M \rightarrow j_*M).$$

The image is taken in the abelian category $\mathbf{MM}(S)$, using the exactness of $j_!$ and j_* , Axiom 4.2.

Remark 5.4. Let $i : Z \rightarrow S$ be the complement of j . The localization triangles (Axiom 1.10) and cohomological amplitude of i^* (Axiom 4.2) yield short exact sequences in $\mathbf{MM}(S)$

$$0 \rightarrow i_* {}^p\text{H}^{-1} i^* j_* M \rightarrow j_! M \rightarrow j_{!*} M \rightarrow 0, \quad (6)$$

$$0 \rightarrow j_{!*} M \rightarrow j_* M \rightarrow i_* {}^p\text{H}^0 i^* j_* M \rightarrow 0. \quad (7)$$

These triangles are the same as for perverse sheaves in the situation that the analog of Axiom 4.2, [BBD82, 4.1.10], is applicable.

Lemma 5.5. • Given any mixed motive M over S' , $j_{!*}M$ is, up to a unique isomorphism, the unique mixed extension of M (i.e., an object X in $\mathbf{MM}(S)$ such that $j^*X = M$) not having nonzero subobjects or quotients of the form i_*N , where $i : Z \rightarrow S$ is the closed complement of j and N is a mixed motive on Z .

- For any two composable open immersions j_1 and j_2 we have $j_{1!*} \circ j_{2!*} = (j_1 \circ j_2)_{!*}$.
- $j_{!*}$ commutes with duals, i.e., $D(j_{!*}-) \cong j_{!*}D(-)$.

Proof: The proofs of the same facts for perverse sheaves [BBD82, Cor. 1.4.25, 2.1.7.1] carry over to this setting. The first statement easily implies the last one. $\square$

Lemma 5.6. (compare [BBD82, Cor. 5.3.2]) The intermediate extension $j_{!*}$ preserves weights of pure objects: given a pure motive $M \in \mathbf{MM}(S')$, $j_{!*}M$ is pure of the same weight.

Proof: This is a consequence of the triangles (6), (7) and the strictness of the weight filtration. $\square$

The following proposition makes precise the intuition that any motive M over S should be reconstructed by its generic fiber (over F) and a finite number of special fibers (over various $\mathbb{F}_p$).

Proposition 5.7. As a thick subcategory of $\mathbf{DM}(S)$, $\mathbf{DM}_{\text{gm}}(S)$ is generated by motives of the form

- $i_*M(X_p)(m)$, where $X_p/\mathbb{F}_p$ is smooth projective, $m \in \mathbb{Z}$ and $i : \text{Spec } \mathbb{F}_p \rightarrow S$ is any closed point and
- $j_{!*}j^*h^k(X, m)$, where X is regular, flat and projective over S with smooth generic fiber, and $j : S' \rightarrow S$ is such that $X \times_S S'$ is smooth over S' and k and m are arbitrary.

Proof: Let $\mathcal{D} \subset \mathbf{DM}_{\text{gm}}(S)$ be the thick category generated by the objects in the statement. By resolution of singularities over S (Axiom 1.15), $\mathbf{DM}_{\text{gm}}(S)$ is the thick subcategory of $\mathbf{DM}(S)$ generated by objects $i_*M(X_p)(m)$ and $M(X)(m)$, where X_p and X are as in the statement and $m \in \mathbb{Z}$.

It is therefore sufficient to see $M := M(X) \in \mathcal{D}$. Let $j : S' \rightarrow S$ be such that $X_{S'}$ is smooth over S' . By 1.10 it is enough to show $j_*j^*M \in \mathcal{D}$, since motives over finite fields are already covered. Applying the truncations with respect to the motivic t -structure to j_*j^*M and exactness of j_* , j^* (Axiom 4.2) shows that we may deal with $j_*j^*h^k(X, m)$ for all k instead of j_*j^*M . (Only finitely many k yield a nonzero term by Axiom 4.1.) By Remark 5.4, there is a short exact sequence of mixed motives

$$0 \rightarrow j_{!*}j^*h^k(X, m) \rightarrow j_*j^*h^k(X, m) \rightarrow i_*{}^p\mathcal{H}^0 i^* j_*j^*h^k(X, m) \rightarrow 0.$$

Here i is the complement of j . The left and right hand motives are in $\mathcal{D}$, hence so is the middle one. $\square$

5.3 Smooth motives

The notion of smooth motives (a neologism leaning on lisse sheaves) is a technical stepstone for the definition of the generic intermediate extension $\eta_{!*}$, cf. Definition 5.14. Roughly speaking, smoothness for mixed motives M means that i^*M and $i^!M$ do not intermingle in the sense that their cohomological degrees are disjoint.

Definition 5.8. Let M be a geometric motive over S . It is called *smooth* if for any closed point $i : \text{Spec } \mathbb{F}_p \rightarrow S$ there is an isomorphism

$$i^!M \cong i^*M(-1)[-2].$$

M is called *generically smooth* if there is an open (non-empty) immersion $j : S' \rightarrow S$ such that j^*M is smooth.

Let X/S be a scheme with smooth generic fiber X_η . Then $M_S(X)$ is a generically smooth motive.

The isomorphism in Definition 5.8 is not required to be canonical in any sense. Therefore, the subcategory of smooth motives is *not* triangulated in $\mathbf{DM}_{\text{gm}}(S)$.

Lemma 5.9. *Let M be a smooth mixed motive over S . Let $i : Z \rightarrow S$ be proper closed subscheme, let $j : S' \rightarrow S$ be its complement. Then $i^!M = (\text{pH}^1 i^!M)[-1]$ and dually $i^*M = (\text{pH}^{-1} i^*M)[1]$.*

Proof: By assumption $i^!M \cong i^*M(-1)[-2]$. By Axiom 4.2, the left hand side of that isomorphism is concentrated in degrees $[0, 1]$. The right hand side is in degrees $[1, 2]$. This shows $i^!M = \text{pH}^1(i^!M)[-1]$ by Axiom 4.1 and similarly for i^*M . $\square$

The following is the key relation of smooth motives and the intermediate extension. Note the similarity with Lemma 5.15.

Lemma 5.10. *Let M be a smooth mixed motive over S . Then M is canonically isomorphic to $j_{!*}j^*M$.*

Proof: Let $i : Z \rightarrow S$ be the complement of j . Given any $i_*N \subset M$ with $N \in \mathbf{MM}(Z)$, we apply the left-exact functor $i^!$ and see $N \subset \text{pH}^0(i^!M) \stackrel{5.9}{=} 0$. Quotients of M of the form i_*N are treated dually. We now invoke Lemma 5.5. $\square$

Lemma 5.11. *Let X be any smooth projective scheme over S . Set $M := M(X)$. Then all $\text{h}^n X = \text{pH}^n M$ are smooth.*

Proof: Let $f_{m,n}$ be the (m, n) -component of the bottom isomorphism making the following commutative:

$$\begin{array}{ccc} i^!M & \xrightarrow{\cong, \text{ see (3) }} & i^*M(-1)[-2] \\ \downarrow \cong, 4.4 & & \downarrow \cong, 4.4 \\ \oplus_m A_m := \oplus i^!(\text{pH}^m M)[-m] & \xrightarrow{\cong} & \oplus_n B_n := \oplus i^*(\text{pH}^n M)(-1)[-n-2]. \end{array}$$

We claim $f_{m,n} = 0$ for all $m \neq n$, from which the lemma follows. By Axiom 4.5 we have $B_n \cong h^{n-1}(X_{\mathbb{P}})[-n-1](-1)$. Using this and the reflexivity of the Verdier dual functor, we obtain an isomorphism $A_m \cong (\mathrm{PH}^{m+1}i^!M)[-1-m]$. Hence B_n is concentrated in cohomological degree $n+1$, while A_m is in degree $n+2$. (The a priori bounds of Axiom 4.2 would be $[m, m+1]$ and $[n+1, n+2]$, respectively.) As the cohomological dimension of motives over $\mathbb{F}_{\mathbb{P}}$ is zero (Axiom 4.1), the only way for $f_{m,n} \neq 0$ is $m = n$. $\square$

5.4 Generic intermediate extension

Lemma 5.12. (*Spreading out morphisms*) *Given two geometric motives M and M' over S together with a map $\phi_{\eta} : \eta^*M \rightarrow \eta^*M'$, there is an open subscheme $j : S' \subset S$ and a map $\phi_{S'} : j^*M \rightarrow j^*M'$ which extends ϕ_{η} . Any two such extensions agree when restricted to a possibly smaller open subscheme. In particular, if ϕ_{η} is an isomorphism, then $\phi_{S'}$ is an isomorphism for sufficiently small S' .*

Proof: The adjunction map $M \rightarrow \eta_*\eta^*M$ and $\eta_*\phi_{\eta}$ give a map $M \rightarrow \eta_*\eta^*M'$, hence by (4) a map $M \rightarrow \bigoplus_{\mathbb{P}} i_{\mathbb{P}*} i_{\mathbb{P}}^! M'[1]$. As M is compact, it factors over a finite sum $\bigoplus_{\mathbb{P} \in T} i_{\mathbb{P}*} i_{\mathbb{P}}^! M'[1]$. Let $j : S' \rightarrow S$ be the complement of the points in T . The map $M \rightarrow \eta_*\eta^*M'$ factors over j_*j^*M' and gives a map $j^*M \rightarrow j^*M'$ which extends ϕ_{η} . The first claim is shown.

For the unicity of the extension, we may assume that ϕ_{η} is zero, and show that $\phi_{S'}$ is zero for some suitable S' . This is the same argument as before: the map $M \rightarrow j_*j^*M'$ constructed in the previous step factors over $\bigoplus_{\mathbb{P} \in S'} i_{\mathbb{P}*} i_{\mathbb{P}}^! M'$, since $M \rightarrow \eta_*\eta^*M'$ is zero. By compactness of M , only finitely many primes in the sum contribute to the map, discarding these yields the claim.

If ϕ_{η} is an isomorphism, $\psi_{\eta} := \phi_{\eta}^{-1}$ can be extended to some $\psi_{S'}$. As both $\phi_{S'} \circ \psi_{S'}$ and $\mathrm{id}_{S'}$ extend id_{η} , they agree on some possibly smaller open subscheme of S and similarly with $\psi_{S'} \circ \phi_{S'}$. $\square$

Remark 5.13. The lemma shows the full faithfulness of the functor

$$\varinjlim_{S' \subset S} \mathbf{DM}_{\mathrm{gm}}(S') \xrightarrow{\eta^*} \mathbf{DM}_{\mathrm{gm}}(F).$$

Its essential surjectivity is a consequence of Axiom 1.5, so we have an equivalence. However, we will stick to the more basic language of colimits in $\mathbf{DM}(S)$ instead of colimits of the categories of geometric motives.

Definition 5.14. Let $M_{\eta} \in \mathbf{DM}_{\mathrm{gm}}(F)$ be a motive such that there exists a generically smooth mixed motive M over S (Definition 5.8) with $\eta^*M \cong M_{\eta}$. Then the *generic intermediate extension* $\eta_{!*}M_{\eta}$ is defined as

$$\eta_{!*}M_{\eta} := j_{!*}j^*M$$

where $j : S' \rightarrow S$ is an open immersion such that j^*M is smooth.

This is independent of the choices of j and M (Lemmas 5.10, 5.12) and functorial (5.12). For a mixed, non-smooth motive M , there need not be a map $j_{!*}j^*M \rightarrow M$. Therefore, $\varinjlim j_{!*}j^*M$ does not make sense unless there is some smoothness constraint on M_η .

5.5 Intermediate extension and ℓ -adic realization

This subsection deals with the interplay of the (generic) intermediate extension functor on mixed motives and the ℓ -adic realization. In this subsection, S is an open subscheme of $\mathrm{Spec} \mathcal{O}_F[1/\ell]$. The following lemma is well-known.

Lemma 5.15. *Let $\mathcal{F}$ be an étale (honest) locally constant sheaf on S . Let $\eta : \mathrm{Spec} F \rightarrow \mathrm{Spec} \mathcal{O}_F[1/\ell]$ be the generic point. Then the canonical map $\mathcal{F} \rightarrow R^0\eta_*\eta^*\mathcal{F}$ is an isomorphism.*

Lemma 5.16. *Let M be a mixed motive over S' . Let $j : S' \rightarrow S$ be an open immersion. Then*

$$(j_{!*}M)_\ell = j_{!*}(M_\ell).$$

Let i be the complementary closed immersion to $j : S' \rightarrow S$ and let η' and η be the generic point of S' and S , respectively. If M is additionally smooth, one has

$$(i^*j_{!*}M)_\ell = i^*j_{!*}M_\ell = i^*(R^0\eta_*\eta'^*M_\ell[-1])[1].$$

To clarify the statement, note that the ℓ -adic realization of M is a perverse sheaf on S' by Axiom 4.8. Thus, $j_{!*}$ (Section 3) can be applied to it.

Proof: The first statement follows from Axiom 2.1, the definition of $j_{!*}$ and the exactness of $R\Gamma_\ell$ (Axiom 4.8).

Let now M be mixed and smooth over S' . As M_ℓ is a perverse sheaf by 4.8, there is an open immersion $j' : S'' \rightarrow S'$ such that $j'^*M_\ell[-1]$ is a locally constant (honest) sheaf on S'' . As M is smooth we know from Lemmas 5.5 and 5.10

$$i^*j_{!*}M = i^*(j \circ j')_{!*}j'^*M.$$

By the interpretation of $(j \circ j')_{!*}$ in terms of $R^0(j \circ j')_*$ (Section 3) we have

$$(i^*j_{!*}M)_\ell = i^*j_{!*}M_\ell = i^*(R^0(j \circ j')_*j'^*M_\ell[-1])[1] \stackrel{5.15}{=} i^*(R^0\eta_*\eta'^*M_\ell[-1])[1].$$

□

6 f -cohomology

6.1 f -cohomology via non-ramification

For any place $\mathfrak{p}$ of F , let $F_{\mathfrak{p}}$ be the completion, $G_{\mathfrak{p}}$ the local Galois group. For finite places, $I_{\mathfrak{p}}$ denotes the inertia group. For brevity, we will usually write $H^*(M)$ for $H^*(S, M)$, where M is any motive over some base S .

Definition 6.1. [BK90, Section 3] Let V be a finite-dimensional ℓ -adic vector space, endowed with a continuous action of $G_{\mathfrak{p}}$, where $\mathfrak{p}$ is a finite place of F not over ℓ . Set

$$H_f^i(F_{\mathfrak{p}}, V) := \begin{cases} H^0(F_{\mathfrak{p}}, V) & i = 0 \\ \ker H^1(F_{\mathfrak{p}}, V) \rightarrow H^1(I_{\mathfrak{p}}, V) & i = 1 \\ 0 & \text{else.} \end{cases}$$

Remark 6.2. If $\mathfrak{p}$ lies over ℓ , the definition is completed by $H_f^1(F_{\mathfrak{p}}, V) := \ker H^1(F_{\mathfrak{p}}, V) \rightarrow H^1(F_{\mathfrak{p}}, B_{crys} \otimes V)$, where B_{crys} denotes the ring of $\mathfrak{p}$ -adic periods [FM87]. We will disregard this case throughout.

Lemma 6.3. Let $\eta_{\mathfrak{p}} : \text{Spec } F_{\mathfrak{p}} \rightarrow \text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}$ be the generic point of the completion of $\mathcal{O}_F$ at $\mathfrak{p}$. Using the above notation, for $\mathfrak{p}$ not over ℓ , there is a canonical isomorphism $H_f^1(F_{\mathfrak{p}}, V) \cong H^1(\mathcal{O}_{F_{\mathfrak{p}}}, R^0 \eta_{\mathfrak{p}*} V)$. (The right hand side denotes ℓ -adic cohomology over $\mathcal{O}_{F_{\mathfrak{p}}}$.)

Proof: Let U_n be an ℓ^n -torsion sheaf on $F_{\mathfrak{p}}$. In this proof, we will write

$$H_f^1(F_{\mathfrak{p}}, U_n) := \ker H^1(F_{\mathfrak{p}}, U_n) \rightarrow H^1(I_{\mathfrak{p}}, U_n),$$

i.e., the same definition as for $\mathbb{Q}_{\ell}$ -sheaves is done. The $\mathbb{Q}_{\ell}$ -sheaf V is, by definition, of the form $U \otimes_{\mathbb{Z}_{\ell}} \mathbb{Q}_{\ell}$, where $U = (U_n)_n$ is a projective system of $\mathbb{Z}/\ell^n$ -sheaves. By definition

$$H^1(F_{\mathfrak{p}}, V) = \varprojlim_{n \in \mathbb{N}} H^1(F_{\mathfrak{p}}, U_n) \otimes \mathbb{Q}_{\ell}$$

and similarly for $H^1(I_{\mathfrak{p}}, V)$. Both $\varprojlim_n$ and $-\otimes_{\mathbb{Z}_{\ell}} \mathbb{Q}_{\ell}$ are left-exact functors, so one has

$$H_f^1(F_{\mathfrak{p}}, V) = \left(\varprojlim_n H_f^1(F_{\mathfrak{p}}, U_n) \right) \otimes \mathbb{Q}_{\ell}.$$

Thus we may henceforth assume that V is a $\mathbb{Z}/\ell^n$ -sheaf.

Recall the description of étale sheaves on $\mathcal{O}_{F_{\mathfrak{p}}}$ from [Mil80, II.3.12, II.3.16]. Let $i : \text{Spec } \mathbb{F}_{\mathfrak{p}} \rightarrow \text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}$ be the closed point. As $\mathcal{O}_{F_{\mathfrak{p}}}$ is a henselian ring [Mil80, Prop. I.4.5], for any sheaf $\mathcal{F}$ on $\text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}$, the global sections depend only on the special fiber and

$$\Gamma_{\text{Spec } F_{\mathfrak{p}}} = \Gamma_{\text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}} \circ (\eta_{\mathfrak{p}*}) = \Gamma_{\text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}} \circ (i_* i^* \eta_{\mathfrak{p}*}).$$

These functors can be interpreted using group cohomology: $\Gamma_{\text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}} \circ i_* = \Gamma_{\mathbb{F}_{\mathfrak{p}}}$ and $(-)^{I_{\mathfrak{p}}} = i^* \eta_{\mathfrak{p}*}$ (*loc. cit.*). The Hochschild-Serre spectral sequence for $(-)^{G_{\mathfrak{p}}} = (-)^{\text{Gal}(\mathbb{F}_{\mathfrak{p}})} \circ (-)^{I_{\mathfrak{p}}}$ can be translated to

$$H^p(\text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}, i_* i^* R^q \eta_{\mathfrak{p}*} V) \Rightarrow H^n(F_{\mathfrak{p}}, V).$$

In addition we have the Leray spectral sequence

$$H^p(\text{Spec } \mathcal{O}_{F_{\mathfrak{p}}}, R^q \eta_{\mathfrak{p}*} V) \Rightarrow H^n(F_{\mathfrak{p}}, V).$$

The exact sequence of low degrees of the Hochschild-Serre sequence maps to the sequence below:

$$\begin{array}{ccccccc}
0 & \longrightarrow & H^1(\mathrm{Spec} \mathcal{O}_{F_{\mathfrak{p}}}, R^0 \eta_{\mathfrak{p}*} V) & \longrightarrow & H^1(F_{\mathfrak{p}}, V) & \longrightarrow & H^0(\mathrm{Spec} \mathcal{O}_{F_{\mathfrak{p}}}, R^1 \eta_{\mathfrak{p}*} V) \\
& & & & \downarrow = & & \downarrow \\
0 & \longrightarrow & H_f^1(F_{\mathfrak{p}}, V) & \longrightarrow & H^1(F_{\mathfrak{p}}, V) & \longrightarrow & H^1(I_{\mathfrak{p}}, V)
\end{array}$$

As $H^0(\mathrm{Gal}(\mathbb{F}_{\mathfrak{p}}), H^1(I_{\mathfrak{p}}, V)) \subset H^1(I_{\mathfrak{p}}, V)$ and $\Gamma_{\mathcal{O}_{F_{\mathfrak{p}}}} = \Gamma_{\mathcal{O}_{F_{\mathfrak{p}}}} \circ i_* i^*$, the right hand map is injective, therefore there is a unique isomorphism between the left hand terms making the diagram commutative. $\square$

In order to proceed to a global level, the following definition is done:

Definition 6.4. [FPR94, II.1.3] Given an ℓ -adic continuous representation V of $G = \mathrm{Gal}(F)$, define $H_f^i(F, V)$ to be such that the following diagram is cartesian. In the lower row, V is considered a $G_{\mathfrak{p}} = \mathrm{Gal}(F_{\mathfrak{p}})$ -module by restriction.

$$\begin{array}{ccc}
H_f^i(F, V) & \longrightarrow & H^i(F, V) \\
\downarrow & & \downarrow \\
\prod H_f^i(F_{\mathfrak{p}}, V) & \longrightarrow & \prod H^i(F_{\mathfrak{p}}, V)
\end{array}$$

The product ranges over all finite places $\mathfrak{p}$ of F . We define $H_{f, \setminus \mathrm{crys}}^i(F, V)$ similarly, except that in the lower row of the preceding diagram only places $\mathfrak{p}$ that do not lie over ℓ occur.

Lemma 6.5. *Let V be an ℓ -adic étale sheaf on $\mathrm{Spec} F$. Then there is a natural isomorphism*

$$H_{f, \setminus \mathrm{crys}}^i(F, V) \cong H^1(\mathcal{O}_F[1/\ell], R^0 \eta_* V).$$

Proof: By the same argument as in the previous proof, we may assume that V is a sheaf of $\mathbb{Z}/\ell^n$ -modules, since the isomorphism we are going to establish is natural in V and

$$H_{f, \setminus \mathrm{crys}}^i(F, V) = \ker H^i(F, V) \rightarrow \prod_{\mathfrak{p} \nmid \ell} (H^i(F_{\mathfrak{p}}, V)/H_f^i(F_{\mathfrak{p}}, V)).$$

Consider the following cartesian diagram ($\mathfrak{p} \nmid \ell$)

$$\begin{array}{ccccc}
\mathrm{Spec} F_{\mathfrak{p}} & \xrightarrow{\eta_{\mathfrak{p}}} & \mathrm{Spec} \mathcal{O}_{F_{\mathfrak{p}}} & \xleftarrow{i_{\mathfrak{p}}} & \mathrm{Spec} \mathbb{F}_{\mathfrak{p}} \\
\downarrow b & & \downarrow a & & \downarrow = \\
\mathrm{Spec} F & \xrightarrow{\eta} & \mathrm{Spec} \mathcal{O}_F[1/\ell] & \xleftarrow{i} & \mathrm{Spec} \mathbb{F}_{\mathfrak{p}}
\end{array}$$

In the derived category of $\mathbb{Z}/\ell^n$ -sheaves on $\mathrm{Spec} \mathcal{O}_F[1/\ell]$, there is a triangle $R^0 \eta_* V \rightarrow R\eta_* V \rightarrow R^1 \eta_*[-1]V$. Likewise, $R^0 \eta_{\mathfrak{p}*} b^* V \rightarrow R\eta_{\mathfrak{p}*} b^* V \rightarrow$

$R^1\eta_{\mathfrak{p}*}b^*V[-1]$. (We have used $\mathfrak{p} \nmid \ell$, since the inertia group has cohomological dimension bigger than one for $\mathfrak{p}|\ell$.) This yields exact horizontal sequences, the vertical maps are adjunction maps

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^1(\mathrm{Spec} \mathcal{O}_F[1/\ell], \eta_* V) & \longrightarrow & H^1(F, V) & \longrightarrow & H^0(\mathrm{Spec} \mathcal{O}_F[1/\ell], R^1\eta_* V) \\ & & \downarrow & & \downarrow & & \downarrow \alpha \\ 0 & \longrightarrow & \prod_{\mathfrak{p} \nmid \ell} H^1(\mathcal{O}_{F_{\mathfrak{p}}}, R^0\eta_{\mathfrak{p}*}b^*V) & \longrightarrow & \prod_{\mathfrak{p} \nmid \ell} H^1(F_{\mathfrak{p}}, b^*V) & \longrightarrow & \prod_{\mathfrak{p} \nmid \ell} H^0(\mathcal{O}_{F_{\mathfrak{p}}}, R^1\eta_{\mathfrak{p}*}b^*V) \end{array}$$

We will show that α is injective. Hence, the left square is cartesian and by definition and Lemma 6.3 the claim is shown. Indeed, α factors as

$$H^0(\mathcal{O}_F[1/\ell], R^1\eta_* V) \subset \prod_{\mathfrak{p} \nmid \ell} H^0(\mathbb{F}_{\mathfrak{p}}, i_{\mathfrak{p}}^* R^1\eta_* V) \rightarrow \prod_{\mathfrak{p} \nmid \ell} H^0(\mathcal{O}_{F_{\mathfrak{p}}}, R^1\eta_{\mathfrak{p}*}b^*V) \left(\cong \prod_{\mathfrak{p} \nmid \ell} H^0(\mathbb{F}_{\mathfrak{p}}, i_{\mathfrak{p}}^* R^1\eta_{\mathfrak{p}*}b^*V) \right).$$

using $i^* R^1\eta_* V = i_{\mathfrak{p}}^* a^* R^1\eta_* V = i_{\mathfrak{p}}^* R^1\eta_{\mathfrak{p}*}b^*V$ (the latter isomorphism is found e.g. in [Mil08, Example 12.4]). $\square$

Definition 6.6. [Beĭ87a, Remark 4.0.1.b], [BK90, Conj. 5.3], [Fon92, Section 6.5], [FPR94, III.3.1.3] Let M_{η} be a mixed motive over F . Let, similarly to Definition 6.4, $H_f^i(M_{\eta})$ be defined such that the following diagram, in which the bottom products are taken over all primes ℓ , is cartesian. As usual, $M_{\eta_{\ell}}$ is the ℓ -adic realization, seen as a G -module.

$$\begin{array}{ccc} H_f^i(F, M_{\eta}) & \longrightarrow & H^i(F, M_{\eta}) \\ \downarrow & & \downarrow \\ \prod_{\ell} H_f^i(F, M_{\eta_{\ell}}) & \longrightarrow & \prod_{\ell} H^i(F, M_{\eta_{\ell}}) \end{array}$$

Again, to rid ourselves from crystalline questions at $\mathfrak{p}|\ell$, we define $H_{f, \setminus \mathrm{crys}}^i(F, M_{\eta})$ by replacing $\prod_{\ell} H_f^i(F, M_{\eta_{\ell}})$ in the bottom row by $\prod_{\ell} H_{f, \setminus \mathrm{crys}}^i(F, M_{\eta_{\ell}})$.

We are now going to exhibit an interpretation of f -cohomology thus defined in terms of the generic intermediate extension η_* . Recall that we are assuming in this section the axioms of Sections 1, 2, and 4. Mixed motives are needed to even define η_* . Moreover, for the comparison result, we need to assume the following conjecture.

Conjecture 6.7. (Beilinson) Let $X/\mathbb{F}_q$ be smooth and projective. Up to torsion, numerical and rational equivalence agree on X .

As rational and numerical equivalence are the finest and the coarsest adequate equivalence relation, respectively [And04, Section 3.2], this conjecture implies that all adequate equivalence relations, in particular homological equivalence agree.

Lemma 6.8. *Let N be any mixed motive over $\mathbb{F}_p$. Conjecture 6.7 implies that the ℓ -adic realization map $H^0(\mathbb{F}_p, N) \rightarrow H^0(\mathbb{F}_p, N_\ell) := N_\ell^{\text{Gal}(\mathbb{F}_p)}$ is injective.*

Proof: We can assume that N is pure of weight 0, since both depend only on $\text{gr}_0^W N$ (Axiom 4.9). Under Conjecture 6.7 all adequate equivalence relations agree, so that we may regard N as a pure motive with respect to any adequate equivalence relation. As the injectivity is stable under taking direct summands, we may assume $N = h(X, n)$ for X smooth and projective over $\mathbb{F}_p$, by definition of pure motives and Axiom 4.13. The left hand side is given by $\text{CH}^n(X)$, so the map is injective by Conjecture 6.7. $\square$

Theorem 6.9. *Let M be a generically smooth mixed motive over $\mathcal{O}_F$ (Definition 5.8). Set $\eta^* M[-1] =: M_\eta$. Assuming Conjecture 6.7, there is a natural isomorphism*

$$H^0(\mathcal{O}_F, \eta_{!*} \eta^* M) \xrightarrow{\cong} H_{f, \setminus \text{crys}}^1(F, M_\eta).$$

Proof: Notice that $\eta_{!*} \eta^* M$ is well-defined by the assumptions. We want to show that there is a cartesian commutative diagram

$$\begin{array}{ccc} H^0(\eta_{!*} \eta^* M) & \longrightarrow & H^0(\eta_* \eta^* M) = H^1(M_\eta) \\ \vdots \downarrow & & \downarrow \\ \prod_\ell H_{f, \setminus \ell}^1(F, M_{\eta_\ell}) & \longrightarrow & \prod_\ell H^1(F, M_{\eta_\ell}) \end{array}$$

Let $j : U \rightarrow \text{Spec } \mathcal{O}_F$ be any open immersion such that $j^* M$ is smooth. We have $\eta_{!*} \eta^* M = j_{!*} j^* M$. The left hand term of the exact sequence

$$\oplus_{\mathfrak{p} \in U} H^0(i_{\mathfrak{p}*} i_{\mathfrak{p}}^! M) \rightarrow H^0(j_* j^* M) \rightarrow H^0(\eta_* \eta^* M) \rightarrow \oplus_{\mathfrak{p} \in U} H^1(i_{\mathfrak{p}*} i_{\mathfrak{p}}^! M)$$

induced by (4) vanishes as $i_{\mathfrak{p}}^! M$ is concentrated in cohomological degree 1 for $\mathfrak{p} \in U$ (Lemma 5.9). Any $a \in H^0(\eta^* M)$ maps to a finite sub-sum of $\oplus_{\mathfrak{p} \in \text{Spec } \mathcal{O}_F} H^1(i_{\mathfrak{p}*} i_{\mathfrak{p}}^! M)$, so letting j be the open complement of these points, a lies in (the image of) $H^0(j_* j^* M)$:

$$H^0(\eta^* M) = \varinjlim_{\substack{j: U \rightarrow \text{Spec } \mathcal{O}_F \\ j^* M \text{ smooth}}} H^0(j_* j^* M).$$

By Lemma 6.10 below, the map $H^0(j_{!*} j^* M) \rightarrow H^0(j_* j^* M) \rightarrow H^0(\eta^* M)$ is injective. Therefore, taking the colimit over all U such that $M|_U$ is smooth, the exact localization sequence

$$0 \rightarrow H^0(j_{!*} j^* M) \rightarrow H^0(j_* j^* M) \rightarrow \oplus_{\mathfrak{p} \notin U} H^0(\text{PH}^0 i_{\mathfrak{p}*} j^* M)$$

stemming from (7) gives

$$0 \rightarrow H^0(j_{!*} j^* M) \rightarrow H^0(\eta_* \eta^* M) \rightarrow \oplus_{\mathfrak{p}} H^0(\text{PH}^0 i_{\mathfrak{p}*} j_{\mathfrak{p}*} j_{\mathfrak{p}}^* M).$$

Here $j_{\mathfrak{p}}$ is the complementary open immersion to $i_{\mathfrak{p}}$ and the direct sum is over all (finite) places $\mathfrak{p}$ of $\mathcal{O}_F$. We have $i_{\mathfrak{p}}^* \eta_* \eta^* M = i_{\mathfrak{p}}^* j_{\mathfrak{p}*} j_{\mathfrak{p}}^* M$, so the top sequence in the following commutative diagram is exact:

$$\begin{array}{ccccccc}
0 & \longrightarrow & H^0(j_{!*} j^* M) & \longrightarrow & H^1(M_{\eta}) & \longrightarrow & \bigoplus_{\mathfrak{p}} H^0({}^p H^0 i_{\mathfrak{p}}^* \eta_* \eta^* M) \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \prod_{\ell} H^0((j_{\ell}^* j_{!*} j^* M)_{\ell}) & \longrightarrow & \prod_{\ell} H^1(M_{\eta_{\ell}}) & \longrightarrow & \prod_{\ell} \prod_{\mathfrak{p} \nmid \ell} H^0({}^p H^0 i_{\mathfrak{p}}^* \eta_* \eta^* M)_{\ell}
\end{array} \tag{8}$$

The lower row denotes ℓ -adic cohomology over $\mathcal{O}_F[1/\ell]$, F , and the various $\mathbb{F}_{\mathfrak{p}}$, respectively. Moreover, $j_{\ell} : \text{Spec } \mathcal{O}_F[1/\ell] \rightarrow \text{Spec } \mathcal{O}_F$ is the open immersion. The remainder of the proof consists in the following steps: we show that the diagram is commutative, that the second row is exact, identify its lower leftmost term and show that the rightmost vertical map is injective. This implies that the left square is cartesian, hence the theorem follows.

We write ι and ι_{ℓ} for the open immersions of $U \cap \text{Spec } \mathcal{O}_F[1/\ell]$ into $\text{Spec } \mathcal{O}_F[1/\ell]$ and U , respectively. By Lemma 5.16 and the exactness of j_{ℓ}^* we have

$$(j_{\ell}^* j_{!*} j^* M)_{\ell} = (\iota_{!*} \iota^* j^* M)_{\ell} = \iota_{!*} \iota^* (j^* M)_{\ell}.$$

Thus (8) is commutative since every term at the bottom just involves the ℓ -adic realization of the motive above it, restricted to $\text{Spec } \mathcal{O}_F[1/\ell]$.

The exactness of the bottom row is shown separately for each ℓ , so ℓ is fixed for this argument. By the characterization just mentioned, $\iota_{!*} \iota^* (j^* M)_{\ell}$ does not change when shrinking U , since $j_{!*} j^* M$ is independent of U (as soon as M is smooth over U). On the other hand, by the exactness of the ℓ -adic realization functor (Axiom 4.8) $(j_{\ell}^* M)_{\ell}$ is a perverse sheaf on $\text{Spec } \mathcal{O}_F[1/\ell]$, so is a locally constant sheaf (shifted into degree -1) on a suitable small open subscheme. Hence we may assume that $\iota^* (j_{\ell}^* M)_{\ell}$ is a locally constant sheaf in degree -1 . By Section 3, $\iota_{!*} \iota^* (j_{\ell}^* M)_{\ell} = (R^0 \iota_{!*} \iota^* (j_{\ell}^* M)_{\ell}[-1])[+1]$, so the lower row is the exact sequence belonging to the distinguished triangle of sheaves on $\text{Spec } \mathcal{O}_F[1/\ell]$

$$R^0 \eta_{\ell*} (M_{\eta})_{\ell} \rightarrow R \eta_{\ell*} (M_{\eta})_{\ell} \rightarrow (R^1 \eta_{\ell*} (M_{\eta})_{\ell})[-1].$$

Here $\eta_{\ell} : \text{Spec } F \rightarrow \text{Spec } \mathcal{O}_F[1/\ell]$ is the generic point. By Lemma 6.5 and Lemma 5.15, the factors in the lower left-hand term of (8) agree with $H_{f, \setminus \text{crys}}^1(F, \eta^* M_{\ell}[-1])$.

To show that the rightmost vertical map of (8) is an injection, let $a = (a_{\mathfrak{p}})_{\mathfrak{p} \in \text{Spec } \mathcal{O}_F}$ be a nonzero element of the rightmost upper term. Only finitely many $a_{\mathfrak{p}}$ are nonzero. Pick some ℓ not lying under any of these indices $\mathfrak{p}$. Then the image of a in $\bigoplus_{\mathfrak{p} \nmid \ell} H^0({}^p H^0 i_{\mathfrak{p}}^* \eta_* \eta^* M)_{\ell} \subset \prod_{\mathfrak{p} \nmid \ell} H^0({}^p H^0 i_{\mathfrak{p}}^* \eta_* \eta^* M)_{\ell}$ is nonzero by Lemma 6.8, which assumes Conjecture 6.7. $\square$

Lemma 6.10. *Let M be a mixed motive over S such that $j^* M$ is smooth for some open immersion $j : U \rightarrow S$. Then both maps $H^0(j_{!*} j^* M) \rightarrow H^0(j^* j^* M) \rightarrow H^0(\eta^* M)$ are injective.*

Proof: Indeed the kernels are $H^{-1}(\mathrm{PH}^0 i^* j_* j^* M) = 0$ and $\bigoplus_{\mathfrak{p} \in U} H^0(i_{\mathfrak{p}}^! M)$, which vanishes since $i_{\mathfrak{p}}^! M$ sits in cohomological degree $+1$, for M is smooth around $\mathfrak{p} \in U$ (Lemma 5.9). $\square$

6.2 f -cohomology via K -theory of regular models

Definition 6.11. Let X_{η} be a smooth and projective variety over F . Let $X/\mathcal{O}_F$ be any projective model $X/\mathcal{O}_F$, i.e., $X \times_{\mathcal{O}_F} F = X_{\eta}$. Then we define

$$H^i(X_{\eta}, n)_{\mathcal{O}_F} := \mathrm{im}(H^i(X, n) \rightarrow H^i(X_{\eta}, n)).$$

Recall that we are assuming the axioms of Sections 1, 2, and 4; the full force of mixed motives will be made use of in the sequel.

Theorem 6.12. *The above is well-defined, i.e., independent of the choice of the model X . More precisely we have natural isomorphisms:*

$$H^0(\eta_* h^{i-1}(X_{\eta}, n)[1]) = \begin{cases} H^i(X_{\eta}, n)_{\mathcal{O}_F} & i < 2n \\ \mathrm{CH}^n(X_{\eta})_{\mathbb{Q}, \mathrm{hom}} & i = 2n \end{cases}$$

Moreover

$$H^{-1}(\eta_* h^{i-1}(X_{\eta}, n)[1]) = H^0(h^{i-1}(X_{\eta}, n)).$$

When X is regular, the definition and the statement are due to Beilinson [Bei86, Lemma 8.3.1]. In this case one has $H^i(X_{\eta}, n)_{\mathcal{O}_F} = \mathrm{im} K'_{2j-i}(X)_{\mathbb{Q}}^{(j)} \rightarrow K'_{2j-i}(X_{\eta})_{\mathbb{Q}}^{(j)}$, but that expression does in general depend on the choice of the model [dJ00, dJ08]. An extension of Beilinson's definition to all Chow motives over F due to Scholl is discussed in the theorem below. We first provide a preparatory lemma.

Lemma 6.13. *Let $M \in \mathrm{MM}(\mathrm{Spec} \mathcal{O}_F)$ be a mixed, generically smooth motive with strictly negative weights (Definition 5.8). Let $j : U \rightarrow \mathrm{Spec} \mathcal{O}_F$ be an open non-empty immersion such that $M|_U$ is smooth. The natural map $j_{!*} j^* M \rightarrow \eta_* \eta^* M$ gives rise to an isomorphism*

$$H^0(j_{!*} j^* M) = \mathrm{im} (H^0(M) \rightarrow H^0(\eta_* \eta^* M)).$$

Proof: By Lemma 6.10, $H^0(j_* j^* M) \rightarrow H^0(\eta_* \eta^* M)$ is injective. Hence it suffices to show $H^0(j_{!*} j^* M) = \mathrm{im}(H^0 M \rightarrow H^0(j_* j^* M))$. Let i be the complement of j . From (6), (7), we have a commutative exact diagram

$$\begin{array}{ccccc} H^0(j_{!*} j^* M) & \xrightarrow{\alpha} & H^0(M) & \longrightarrow & H^0(i_* i^* M) \\ \downarrow & & \downarrow & & \\ 0 = H^{-1}(i_* \mathrm{PH}^0 i^* j_* j^* M) & \longrightarrow & H^0(j_{!*} j^* M) & \longrightarrow & H^0(j_* j^* M) \\ & & \downarrow & & \\ & & H^1(i_* \mathrm{PH}^{-1} i^* j_* j^* M) & = & 0 \end{array}$$

The indicated vanishings are because of t -structure reasons and Axiom 4.1, respectively. It remains to show that α is surjective. As i^*M is concentrated in cohomological degrees $[-1, 0]$ (Axiom 4.2), there is an exact sequence

$$0 = H^1(\mathrm{PH}^{-1}i^*M) \rightarrow H^0(i^*M) \rightarrow H^0(\mathrm{PH}^0i^*M).$$

However $H^0(\mathrm{PH}^0i^*M) = 0$ as i^* preserves negative weights (Axiom 4.11) and by strictness of the weight filtration and compatibility with the t -structure (Axiom 4.9). $\square$

Proof: Let $j : U \rightarrow \mathrm{Spec} \mathcal{O}_F$ be an open nonempty immersion (which exists by smoothness of X_η) such that X_U is smooth over U . By definition of $\eta_{!*}$ and Lemmas 5.1 and 5.11, the left hand term in the theorem agrees with $H^0(j_{!*}h^i(X_U, n))$. In the sequel, we write $M := h^{2n}(X, n)$ and $M_\eta := \eta^*[-1]M = h^{i-1}(X_\eta, n)$.

We first do the case $i < 2n$. The spectral sequences

$$H^a(h^b(X, n)) \Rightarrow H^{a+b}(X, n), \quad H^a(h^b(X_\eta, n)) \Rightarrow H^{a+b}(X_\eta, n)$$

resulting from repeatedly applying truncation functors of the motivic t -structure converge since the cohomological dimension is finite (Axiom 4.1 over F , Lemma 5.2 over $\mathcal{O}_F$). By Lemma 5.2, $H^i(-)$, applied to mixed motives over $\mathcal{O}_F$, is non-zero for $i \in \{-1, 0, 1\}$ only. We thus have to consider two exact sequences. The exact functor $\eta^*[-1]$ maps to similar exact sequences for motivic cohomology over F (the indices work out properly, see Lemma 5.1):

$$\begin{array}{ccccccc} 0 & \longrightarrow & K & \longrightarrow & H^i(X, n) & \longrightarrow & H^{-1}(h^{i+1}(X, n)) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & K_\eta & \longrightarrow & H^i(X_\eta, n) & \longrightarrow & H^0(h^i(X_\eta, n)) \xrightarrow{4.9, 4.12} 0 \longrightarrow 0 \end{array} \quad (9)$$

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^1(h^{i-1}(X, n)) & \longrightarrow & K & \longrightarrow & H^0(M) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H^2(h^{i-2}(X_\eta, n)) \xrightarrow{4.1} 0 & \longrightarrow & K_\eta & \longrightarrow & H^1(M_\eta) \longrightarrow 0 \end{array} \quad (10)$$

Here, K and K_η are certain E_3 -terms of the spectral sequences above. The rightmost vertical map in (9) is injective as one sees by combining (4) with the left-exactness of $i_p^!$. Hence

$$\begin{aligned} H^i(X_\eta, n)_{\mathcal{O}_F} &= \mathrm{im}(H^i(X, n) \rightarrow H^i(X_\eta, n)) = \mathrm{im}(K \rightarrow K_\eta) \\ &= \mathrm{im}(H^0(M) \rightarrow H^1(M_\eta)) \end{aligned}$$

The motive $M = h^i(X, n)$ is a generically smooth (mixed) motive by Lemma 5.11. (Recall that this uses the decomposition axiom 4.5 for smooth projective

varieties.) By Example 4.12, its weights are strictly negative. Thus Lemma 6.13 applies and the case $i < 2n$ is shown.

We now do the case $i = 2n$. Now ${}^p\mathrm{H}^1 i^! j_{!*} j^* M = {}^p\mathrm{H}^1 i^! j_* j^* M$ has strictly positive weights because of the choice of j , Example 4.12, Lemma 5.6, Axiom 4.11, and the compatibility of weights and the motivic t -structure, i.e., $\mathrm{wt} {}^p\mathrm{H}^1(-) \subset \mathrm{wt}(-) + 1$. Therefore $\mathrm{H}^0({}^p\mathrm{H}^1 i^! E) = 0$, $E := j_{!*} j^* M$. Here i is any closed immersion. The localization triangle (4) yields

$$\mathrm{H}^0(E) \xrightarrow{\alpha} \mathrm{H}^0(\eta_* \eta^* E) \stackrel{4.7}{=} \mathrm{CH}^n(X_\eta)_{\mathbb{Q}, \mathrm{hom}} \rightarrow \oplus_{\mathfrak{p}} \mathrm{H}^1(i_{\mathfrak{p}}^! E) = \oplus \mathrm{H}^0({}^p\mathrm{H}^1(i^! E)) = 0.$$

Therefore, α is surjective. The injectivity of α is Lemma 6.10.

To calculate $\mathrm{H}^{-1}(\eta_* M_\eta[1])$, let $j : U \rightarrow \mathrm{Spec} \mathcal{O}_F$ be as above and let i be its complement. The natural map $\mathrm{H}^{-1}(\mathrm{Spec} \mathcal{O}_F, j_{!*} j^* M) \rightarrow \mathrm{H}^{-1}(U, j^* M)$ is an isomorphism by the exact cohomology sequence belonging to (7). Thus we have to show

$$\mathrm{H}^{-1}(\mathrm{Spec} \mathcal{O}_F, j_{!*} j^* M) = \mathrm{H}^{-1}(\mathrm{Spec} \mathcal{O}_F, \eta_* \eta^* M).$$

This follows from the localization axiom 1.10 and $i_{\mathfrak{p}}^! M$ being in cohomological degree $+1$ for all points $\mathfrak{p}$ in U (Lemma 5.9), so that $\mathrm{H}^0(\mathbb{F}_{\mathfrak{p}}, i_{\mathfrak{p}}^! M) = \mathrm{H}^{-1}(\mathbb{F}_{\mathfrak{p}}, i_{\mathfrak{p}}^! M) = 0$. $\square$

By a theorem of Scholl [Sch00, Thm. 1.1.6], there is a unique functorial and additive (i.e., converting finite disjoint unions into direct sums) way to extend the definition of $\mathrm{H}^i(X_\eta, n)_{\mathcal{O}_F}$ as the image of the K -theory of a *regular* proper flat model (Definition 6.11) to all Chow motives over F , in particular to ones of smooth projective varieties X_η/F that do not possess a regular proper model X . The following theorem compares this definition with the one via intermediate extensions.

Theorem 6.14. *Let h_η be a direct summand in the category of Chow motives of $h(X_\eta, n)$ where X_η/F is smooth projective. Let $i \in \mathbb{Z}$ be such that $i - 2n < 0$. Let $\iota : \mathbf{M}_{\mathrm{rat}}(F) \rightarrow \mathbf{DM}_{\mathrm{gm}}(F)$ be the embedding. Then, the group*

$$\mathrm{H}^i(h_\eta)_{\mathcal{O}_F} := \mathrm{H}^0(\eta_*({}^p\mathrm{H}^{i-2n-1}(\iota(h_\eta))[1])).$$

is well-defined and agrees with the aforementioned definition by Scholl.

Proof: Recall $\iota(h(X_\eta, n)) = \mathrm{M}(X_\eta, n)[2n] \in \mathbf{DM}_{\mathrm{gm}}(F)$. We first check that the group is well-defined: let $X/\mathcal{O}_F$ be a projective model of X_η . By Lemma 5.12, there is some model $M \in \mathbf{MM}(\mathcal{O}_F)$ of ${}^p\mathrm{H}^{i-2n-1} \iota(h_\eta)[1]$ and an open subscheme U of $\mathrm{Spec} \mathcal{O}_F$ such that M is a direct summand of ${}^p\mathrm{H}^{i-1} \mathrm{M}(X)(n)$ and such that $X \times U$ is smooth over U . Then $h^{i-1}(X, n)$ is a smooth motive when restricted to U (Lemma 5.11). Hence so is M . Thus $\eta_{!*}$ can be applied to $({}^p\mathrm{H}^{i-2n-1} \iota(h_\eta))[1]$.

The assignment $h_\eta \mapsto \mathrm{H}^0(\eta_{!*}({}^p\mathrm{H}^{i-2n-1} \iota(h_\eta))[1])$ is functorial and additive and $h(X_\eta)(n)$ maps to

$$\mathrm{H}^0(\eta_{!*}({}^p\mathrm{H}^{i-1} \mathrm{M}(X_\eta, n))[1]) \stackrel{6.12}{=} \mathrm{H}^i(X_\eta, n)_{\mathcal{O}_F}.$$

Thus the two definitions agree by Scholl's theorem. $\square$

References

- [And04] Yves André. *Une introduction aux motifs (motifs purs, motifs mixtes, périodes)*, volume 17 of *Panoramas et Synthèses [Panoramas and Syntheses]*. Société Mathématique de France, Paris, 2004.
- [Ayo07] Joseph Ayoub. Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. I. *Astérisque*, (314):x+466 pp. (2008), 2007.
- [BBD82] A. A. Beilinson, J. Bernstein, and P. Deligne. Faisceaux pervers. In *Analysis and topology on singular spaces, I (Luminy, 1981)*, volume 100 of *Astérisque*, pages 5–171. Soc. Math. France, Paris, 1982.
- [Beĭ84] A. A. Beilinson. Higher regulators and values of L -functions. In *Current problems in mathematics, Vol. 24, Itogi Nauki i Tekhniki*, pages 181–238. Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1984.
- [Beĭ86] A. A. Beilinson. Notes on absolute Hodge cohomology. In *Applications of algebraic K -theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983)*, volume 55 of *Contemp. Math.*, pages 35–68. Amer. Math. Soc., Providence, RI, 1986.
- [Beĭ87a] A. A. Beilinson. Height pairing between algebraic cycles. In *K -theory, arithmetic and geometry (Moscow, 1984–1986)*, volume 1289 of *Lecture Notes in Math.*, pages 1–25. Springer, Berlin, 1987.
- [Beĭ87b] A. A. Beilinson. On the derived category of perverse sheaves. In *K -theory, arithmetic and geometry (Moscow, 1984–1986)*, volume 1289 of *Lecture Notes in Math.*, pages 27–41. Springer, Berlin, 1987.
- [BK90] Spencer Bloch and Kazuya Kato. L -functions and Tamagawa numbers of motives. In *The Grothendieck Festschrift, Vol. I*, volume 86 of *Progr. Math.*, pages 333–400. Birkhäuser Boston, Boston, MA, 1990.
- [Bon09] M. V. Bondarko. Differential graded motives: weight complex, weight filtrations and spectral sequences for realizations; Voevodsky versus Hanamura. *J. Inst. Math. Jussieu*, 8(1):39–97, 2009.
- [BV08] Alexander Beilinson and Vadim Vologodsky. A DG guide to Voevodsky’s motives. *Geom. Funct. Anal.*, 17(6):1709–1787, 2008.
- [CD10] Denis-Charles Cisinski and Frédéric Déglise. Triangulated categories of motives. Preprint, 2010.
- [Del74] Pierre Deligne. Théorie de Hodge. III. *Inst. Hautes Études Sci. Publ. Math.*, 44:5–77, 1974.
- [Del80] Pierre Deligne. La conjecture de Weil. II. *Inst. Hautes Études Sci. Publ. Math.*, 52:137–252, 1980.

- [Del94] Pierre Deligne. Décompositions dans la catégorie dérivée. In *Motives (Seattle, WA, 1991)*, volume 55 of *Proc. Sympos. Pure Math.*, pages 115–128. Amer. Math. Soc., Providence, RI, 1994.
- [dJ00] Rob de Jeu. Appendix to the paper of Scholl: a counterexample to a conjecture of Beilinson. In *The arithmetic and geometry of algebraic cycles (Banff, AB, 1998)*, volume 548 of *NATO Sci. Ser. C Math. Phys. Sci.*, pages 491–493. Kluwer Acad. Publ., Dordrecht, 2000.
- [dJ08] Rob de Jeu. Further counterexamples to a conjecture of Beilinson. *J. K-Theory*, 1(1):169–173, 2008.
- [Eke90] Torsten Ekedahl. On the adic formalism. In *The Grothendieck Festschrift, Vol. II*, volume 87 of *Progr. Math.*, pages 197–218. Birkhäuser Boston, Boston, MA, 1990.
- [FM87] Jean-Marc Fontaine and William Messing. p -adic periods and p -adic étale cohomology. In *Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985)*, volume 67 of *Contemp. Math.*, pages 179–207. Amer. Math. Soc., Providence, RI, 1987.
- [Fon92] Jean-Marc Fontaine. Valeurs spéciales des fonctions L des motifs. *Astérisque*, (206):Exp. No. 751, 4, 205–249, 1992. Séminaire Bourbaki, Vol. 1991/92.
- [FPR94] Jean-Marc Fontaine and Bernadette Perrin-Riou. Autour des conjectures de Bloch et Kato: cohomologie galoisienne et valeurs de fonctions L . In *Motives (Seattle, WA, 1991)*, volume 55 of *Proc. Sympos. Pure Math.*, pages 599–706. Amer. Math. Soc., Providence, RI, 1994.
- [Gil80] Henri Gillet. Riemann-Roch theorems for higher algebraic k -theory. *Bull. Amer. Math. Soc. (N.S.)*, 3(2):849–852, 1980.
- [Han95] Masaki Hanamura. Mixed motives and algebraic cycles. I. *Math. Res. Lett.*, 2(6):811–821, 1995.
- [HK06] Annette Huber and Bruno Kahn. The slice filtration and mixed Tate motives. *Compos. Math.*, 142(4):907–936, 2006.
- [Hub95] Annette Huber. *Mixed motives and their realization in derived categories*, volume 1604 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1995.
- [Hub97] Annette Huber. Mixed perverse sheaves for schemes over number fields. *Compositio Math.*, 108(1):107–121, 1997.
- [Hub00] Annette Huber. Realization of Voevodsky’s motives. *J. Algebraic Geom.*, 9(4):755–799, 2000.
- [Ive86] Birger Iversen. *Cohomology of sheaves*. Universitext. Springer-Verlag, Berlin, 1986.

- [Ivo07] Florian Ivorra. Réalisation l -adique des motifs triangulés géométriques. I. *Doc. Math.*, 12:607–671, 2007.
- [KW01] Reinhardt Kiehl and Rainer Weissauer. *Weil conjectures, perverse sheaves and l -adic Fourier transform*, volume 42 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2001.
- [Lev93] Marc Levine. Tate motives and the vanishing conjectures for algebraic K -theory. In *Algebraic K-theory and algebraic topology (Lake Louise, AB, 1991)*, volume 407 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 167–188. Kluwer Acad. Publ., Dordrecht, 1993.
- [Lev98] Marc Levine. *Mixed motives*, volume 57 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1998.
- [Mil80] James S. Milne. *Étale cohomology*, volume 33 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1980.
- [Mil08] James S. Milne. *Lectures on Etale Cohomology*. 2008.
- [Nee01] Amnon Neeman. *Triangulated categories*, volume 148 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001.
- [Org04] Fabrice Orgogozo. Isomotifs de dimension inférieure ou égale à un. *Manuscripta Math.*, 115(3):339–360, 2004.
- [PS08] Chris A. M. Peters and Joseph H. M. Steenbrink. *Mixed Hodge structures*, volume 52 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2008.
- [Qui73] Daniel Quillen. Higher algebraic K -theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [Scha] Jakob Scholbach. Mixed Artin-Tate motives over number rings. Preprint.
- [Schb] Anthony J. Scholl. Integral elements of K -theory and products of modular curves II. Preprint, <http://arxiv.org/abs/0710.5453>.
- [Sch91] Anthony J. Scholl. Remarks on special values of L -functions. In *L -functions and arithmetic (Durham, 1989)*, volume 153 of *London Math. Soc. Lecture Note Ser.*, pages 373–392. Cambridge Univ. Press, Cambridge, 1991.

- [Sch00] Anthony J. Scholl. Integral elements in K -theory and products of modular curves. In *The arithmetic and geometry of algebraic cycles (Banff, AB, 1998)*, volume 548 of *NATO Sci. Ser. C Math. Phys. Sci.*, pages 467–489. Kluwer Acad. Publ., Dordrecht, 2000.
- [Sch10] Jakob Scholbach. Special L -values of geometric motives. 2010.
- [SGA72] *Théorie des topos et cohomologie étale des schémas. Tome 2*. Springer-Verlag, Berlin, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat, Lecture Notes in Mathematics, Vol. 270.
- [SGA73] *Théorie des topos et cohomologie étale des schémas. Tome 3*. Lecture Notes in Mathematics, Vol. 305. Springer-Verlag, Berlin, 1973. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de P. Deligne et B. Saint-Donat.
- [Voe00] Vladimir Voevodsky. Triangulated categories of motives over a field. In *Cycles, transfers, and motivic homology theories*, volume 143 of *Ann. of Math. Stud.*, pages 188–238. Princeton Univ. Press, Princeton, NJ, 2000.
- [Wil] J. Wildeshaus. Notes on Artin-Tate motives. Preprint, Nov 2008, arXiv, <http://arxiv.org/abs/0811.4551v1>.